



Special and General Relativity

Boud Roukema

(c) CC-BY-SA-3.0



GR: intro

1. spacetime = 4D (curved) pseudo-Riemannian manifold M with metric g

GR: intro

1. spacetime = 4D (curved) pseudo-Riemannian manifold M with metric g
2. $\forall$ spacetime point $x \exists$ 4D Minkowski tangent space $T_x M$ at x

GR: intro

1. spacetime = 4D (curved) pseudo-Riemannian manifold M with metric g
2. $\forall$ spacetime point $x \exists$ 4D Minkowski tangent space $T_x M$ at x
= vector space (e.g. 4-momentum vectors)

GR: intro

1. spacetime = 4D (curved) pseudo-Riemannian manifold M with metric g
2. $\forall$ spacetime point $x \exists$ 4D Minkowski tangent space $T_x M$ at x
= vector space, $g \Rightarrow$ lengths of vectors in $T_x M$

GR: intro

1. spacetime = 4D (curved) pseudo-Riemannian manifold M with metric g
2. $\forall$ spacetime point $x \in M \exists$ 4D Minkowski tangent space $T_x M$ at x
= vector space, $g \Rightarrow$ lengths of vectors in $T_x M$
3. also, $\forall x \in M, \exists$ 4D Mink. cotangent space $T_x^* M$

GR: intro

1. spacetime = 4D (curved) pseudo-Riemannian manifold M with metric g
2. $\forall$ spacetime point $x \in M \exists$ 4D Minkowski tangent space $T_x M$ at x
= vector space, $g \Rightarrow$ lengths of vectors in $T_x M$
3. also, $\forall x \in M, \exists$ 4D Mink. cotangent space $T_x^* M$
= dual vector space (think: contour map, gradients)

GR: intro

1. spacetime = 4D (curved) pseudo-Riemannian manifold M with metric g

2. $\forall$ spacetime point $x \in M \exists$ 4D Minkowski tangent space $T_x M$ at x

= vector space, $g \Rightarrow$ lengths of vectors in $T_x M$

3. also, $\forall x \in M, \exists$ 4D Mink. cotangent space $T_x^* M$

= space of one-forms, $g^{-1} \Rightarrow$ “lengths”

GR: intro

1. spacetime = 4D (curved) pseudo-Riemannian manifold M with metric g

2. $\forall$ spacetime point $x \in M$ $\exists$ 4D Minkowski tangent space $T_x M$ at x

= vector space, $g \Rightarrow$ lengths of vectors in $T_x M$

3. also, $\forall x \in M$, $\exists$ 4D Mink. cotangent space $T_x^* M$

= space of one-forms, $g^{-1} \Rightarrow$ “lengths”

duality in a basis of $T_x M$ and a basis of $T_x^* M$ usually defined using δ^μ_ν

GR: intro

1. spacetime = 4D (curved) pseudo-Riemannian manifold M with metric g

2. $\forall$ spacetime point $x \in M \exists$ 4D Minkowski tangent space $T_x M$ at x

= vector space, $g \Rightarrow$ lengths of vectors in $T_x M$

3. also, $\forall x \in M, \exists$ 4D Mink. cotangent space $T_x^* M$

= space of one-forms, $g^{-1} \Rightarrow$ “lengths”

2+3. vector–one-form duality in a basis: δ^μ_ν


GR: intro

1. spacetime = 4D (curved) pseudo-Riemannian manifold M with metric g
2. $\forall$ spacetime point $x \in M \exists$ 4D Minkowski tangent space $T_x M$ at x
= vector space, $g \Rightarrow$ lengths of vectors in $T_x M$
3. also, $\forall x \in M, \exists$ 4D Mink. cotangent space $T_x^* M$
= space of one-forms, $g^{-1} \Rightarrow$ “lengths”
- 2+3. vector–one-form duality in a basis: δ^μ_ν
4. w:Levi-Civita connection $\Leftarrow$ metric


GR: intro

1. spacetime = 4D (curved) pseudo-Riemannian manifold M with metric g
2. $\forall$ spacetime point $x \in M \exists$ 4D Minkowski tangent space $T_x M$ at x
= vector space, $g \Rightarrow$ lengths of vectors in $T_x M$
3. also, $\forall x \in M, \exists$ 4D Mink. cotangent space $T_x^* M$
= space of one-forms, $g^{-1} \Rightarrow$ “lengths”
- 2+3. vector–one-form duality in a basis: δ^μ_ν
4. w:Levi-Civita connection $\Leftarrow$ metric
5. metric $\Leftarrow$ Einstein field equations


GR: coordinate transformations




GR: coordinate transformations



GR: coordinate transformations




GR: coordinate transformations



$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$$




GR: coordinate transformations



$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \Lambda \begin{pmatrix} x \\ y \end{pmatrix}$$



GR: coordinate transformations




but

$$\begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$




GR: coordinate transformations



$$\begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$



GR: coordinate transformations




$$\vec{e}_{x'} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \vec{e}_x + \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \vec{e}_y$$




GR: coordinate transformations

$$\vec{e}_{x'} = \Lambda_{x'}^x \vec{e}_x + \Lambda_{x'}^y \vec{e}_y$$



GR: coordinate transformations




also:

$$\begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$




GR: coordinate transformations



$$\vec{e}_{y'} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \vec{e}_x + \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \vec{e}_y$$



GR: coordinate transformations



summary:

$$\vec{e}_{x'} = \Lambda_{x'}^x \vec{e}_x + \Lambda_{x'}^y \vec{e}_y,$$

$$\vec{e}_{y'} = \Lambda_{y'}^x \vec{e}_x + \Lambda_{y'}^y \vec{e}_y,$$

where $\Lambda_{\beta'}^{\alpha} :=$ element
of inverse of $\Lambda_{\beta}^{\alpha'}$,

$$\begin{pmatrix} x' \\ y' \end{pmatrix} = \Lambda \begin{pmatrix} x \\ y \end{pmatrix}$$

GR: coordinate transformations

$$\begin{aligned}\vec{e}_{x'} &= \Lambda_{x'}^x \vec{e}_x + \Lambda_{x'}^y \vec{e}_y, \\ \vec{e}_{y'} &= \Lambda_{y'}^x \vec{e}_x + \Lambda_{y'}^y \vec{e}_y,\end{aligned}\quad \vec{p} \xrightarrow{\mathcal{O}'} \begin{pmatrix} x' \\ y' \end{pmatrix} = \Lambda \begin{pmatrix} x \\ y \end{pmatrix}$$

GR: coordinate transformations

$$\begin{aligned}\vec{e}_{x'} &= \Lambda_{x'}^x \vec{e}_x + \Lambda_{x'}^y \vec{e}_y, \\ \vec{e}_{y'} &= \Lambda_{y'}^x \vec{e}_x + \Lambda_{y'}^y \vec{e}_y, \\ \vec{p} &\rightarrow_{\mathcal{O}'} \begin{pmatrix} x' \\ y' \end{pmatrix} = \Lambda \begin{pmatrix} x \\ y \end{pmatrix}\end{aligned}$$

$$\vec{p} = \sum_i p^i \vec{e}_i$$

GR: coordinate transformations

$$\begin{aligned}\vec{e}_{x'} &= \Lambda_{x'}^x \vec{e}_x + \Lambda_{x'}^y \vec{e}_y, \\ \vec{e}_{y'} &= \Lambda_{y'}^x \vec{e}_x + \Lambda_{y'}^y \vec{e}_y, \end{aligned} \quad \vec{p} \rightarrow_{\mathcal{O}'} \begin{pmatrix} x' \\ y' \end{pmatrix} = \Lambda \begin{pmatrix} x \\ y \end{pmatrix}$$

$$\vec{p} = p^i \vec{e}_i \text{ (w:Einstein summation)}$$

GR: coordinate transformations

$$\begin{aligned}\vec{e}_{x'} &= \Lambda_{x'}^x \vec{e}_x + \Lambda_{x'}^y \vec{e}_y, \\ \vec{e}_{y'} &= \Lambda_{y'}^x \vec{e}_x + \Lambda_{y'}^y \vec{e}_y, \quad \vec{p} \rightarrow_{\mathcal{O}'} \begin{pmatrix} x' \\ y' \end{pmatrix} = \Lambda \begin{pmatrix} x \\ y \end{pmatrix}\end{aligned}$$

$$\vec{p} = p^i \vec{e}_i \text{ (w:Einstein summation)}$$

Einstein summation:

coordinates like r, θ, x, y :

not a sum: $\Lambda_{y'}^x \vec{e}_x$

repeated up-down coordinate indices like $i, j \in \{0, 1, 2\}$ or $\alpha, \beta, \gamma, \lambda, \mu, \nu \in \{0, 1, 2, 3\}$:

sum: $\Lambda_{j'}^i \vec{e}_i := \Lambda_{y'}^x \vec{e}_x + \Lambda_{y'}^y \vec{e}_y$ for a 2D manifold, coords x, y

GR: coordinate transformations

$$\begin{aligned}\vec{e}_{x'} &= \Lambda_{x'}^x \vec{e}_x + \Lambda_{x'}^y \vec{e}_y, \\ \vec{e}_{y'} &= \Lambda_{y'}^x \vec{e}_x + \Lambda_{y'}^y \vec{e}_y, \end{aligned} \quad \vec{p} \rightarrow_{\mathcal{O}'} \begin{pmatrix} x' \\ y' \end{pmatrix} = \Lambda \begin{pmatrix} x \\ y \end{pmatrix}$$

$$\vec{p} = p^i \vec{e}_i \text{ (w:Einstein summation)}$$

new basis vectors = sum of inverse Λ $\times$ **old** vectors

GR: coordinate transformations

$$\begin{aligned}\vec{e}_{x'} &= \Lambda_{x'}^x \vec{e}_x + \Lambda_{x'}^y \vec{e}_y, \\ \vec{e}_{y'} &= \Lambda_{y'}^x \vec{e}_x + \Lambda_{y'}^y \vec{e}_y, \quad \vec{p} \rightarrow_{\mathcal{O}'} \begin{pmatrix} x' \\ y' \end{pmatrix} = \Lambda \begin{pmatrix} x \\ y \end{pmatrix}\end{aligned}$$

$$\vec{p} = p^i \vec{e}_i \text{ (w:Einstein summation)}$$

new basis vectors = sum of inverse $\Lambda \times$ **old** vectors

$$\vec{e}_{\mu'} = \sum_{\nu} \Lambda_{\mu'}^{\nu} \vec{e}_{\nu}$$

GR: coordinate transformations

$$\begin{aligned}\vec{e}_{x'} &= \Lambda_{x'}^x \vec{e}_x + \Lambda_{x'}^y \vec{e}_y, \\ \vec{e}_{y'} &= \Lambda_{y'}^x \vec{e}_x + \Lambda_{y'}^y \vec{e}_y, \end{aligned} \quad \vec{p} \rightarrow_{\mathcal{O}'} \begin{pmatrix} x' \\ y' \end{pmatrix} = \Lambda \begin{pmatrix} x \\ y \end{pmatrix}$$

$$\vec{p} = p^i \vec{e}_i \text{ (w:Einstein summation)}$$

new basis vectors = sum of inverse Λ $\times$ **old** vectors

$$\vec{e}_{\mu'} = \Lambda_{\mu'}^{\nu} \vec{e}_{\nu}$$

GR: coordinate transformations

$$\begin{aligned}\vec{e}_{x'} &= \Lambda_{x'}^x \vec{e}_x + \Lambda_{x'}^y \vec{e}_y, \\ \vec{e}_{y'} &= \Lambda_{y'}^x \vec{e}_x + \Lambda_{y'}^y \vec{e}_y, \quad \vec{p} \rightarrow_{\mathcal{O}'} \begin{pmatrix} x' \\ y' \end{pmatrix} = \Lambda \begin{pmatrix} x \\ y \end{pmatrix}\end{aligned}$$

$$\vec{p} = p^i \vec{e}_i \text{ (w:Einstein summation)}$$

new basis vectors = sum of inverse $\Lambda \times$ **old** vectors

new coords of vector $\vec{p} = \Lambda \times$ old coords of **same** vector $\vec{p}$

GR: coordinate transformations

$$\begin{aligned}\vec{e}_{x'} &= \Lambda_{x'}^x \vec{e}_x + \Lambda_{x'}^y \vec{e}_y, \\ \vec{e}_{y'} &= \Lambda_{y'}^x \vec{e}_x + \Lambda_{y'}^y \vec{e}_y, \end{aligned} \quad \vec{p} \rightarrow \mathcal{O}' \begin{pmatrix} x' \\ y' \end{pmatrix} = \Lambda \begin{pmatrix} x \\ y \end{pmatrix}$$

$$\vec{p} = p^i \vec{e}_i \text{ (w:Einstein summation)}$$

new basis vectors = sum of inverse Λ $\times$ **old** vectors

vector invariance requires contravariance of its coords

“contra” = inverse of change of basis vectors

GR: coordinate transformations

$$\begin{aligned}\vec{e}_{x'} &= \Lambda_{x'}^x \vec{e}_x + \Lambda_{x'}^y \vec{e}_y, \\ \vec{e}_{y'} &= \Lambda_{y'}^x \vec{e}_x + \Lambda_{y'}^y \vec{e}_y, \end{aligned} \quad \vec{p} \rightarrow \mathcal{O}' \begin{pmatrix} x' \\ y' \end{pmatrix} = \Lambda \begin{pmatrix} x \\ y \end{pmatrix}$$

$$\vec{p} = p^i \vec{e}_i \text{ (w:Einstein summation)}$$

new basis vectors = sum of inverse Λ $\times$ **old** vectors

vector invariance requires contravariance of its coords

“contra” = inverse of change of basis vectors

- $\vec{p}$ is invariant: no dependence on coords
- $\vec{p}$ is contravariant: p^i change inversely to $\vec{e}_i$

GR: coord. transf.: 1-forms

ϕ = scalar field = $\phi(x, y) \equiv \phi(x', y')$

write $\phi_{,x} := \frac{\partial \phi}{\partial x} =: (\tilde{d}\phi)_x$

GR: coord. transf.: 1-forms

ϕ = scalar field = $\phi(x, y) \equiv \phi(x', y')$

write $\phi_{,x} := \frac{\partial \phi}{\partial x} =: (\tilde{d}\phi)_x$

What is the relation between $(\phi_{,x'}, \phi_{,y'})$
and $(\phi_{,x}, \phi_{,y})$?

GR: coord. transf.: 1-forms

ϕ = scalar field = $\phi(x, y) \equiv \phi(x', y')$

write $\phi_{,x} := \frac{\partial \phi}{\partial x} =: (\tilde{d}\phi)_x$

ϕ depends either on x and y , or on x' and y'

$$\Rightarrow \frac{\partial \phi}{\partial x'} = \frac{\partial \phi}{\partial x} \frac{\partial x}{\partial x'} + \frac{\partial \phi}{\partial y} \frac{\partial y}{\partial x'}$$

GR: coord. transf.: 1-forms

ϕ = scalar field = $\phi(x, y) \equiv \phi(x', y')$

write $\phi_{,x} := \frac{\partial \phi}{\partial x} =: (\tilde{d}\phi)_x$

ϕ depends either on x and y , or on x' and y'

$$\Rightarrow \phi_{,x'} = \phi_{,x} x_{,x'} + \phi_{,y} y_{,x'}$$

GR: coord. transf.: 1-forms

ϕ = scalar field = $\phi(x, y) \equiv \phi(x', y')$

write $\phi_{,x} := \frac{\partial \phi}{\partial x} =: (\tilde{d}\phi)_x$

ϕ depends either on x and y , or on x' and y'

$$\Rightarrow \phi_{,x'} = \phi_{,x} x_{,x'} + \phi_{,y} y_{,x'}$$

$$(\phi_{,x'}, \phi_{,y'}) =$$

GR: coord. transf.: 1-forms

ϕ = scalar field = $\phi(x, y) \equiv \phi(x', y')$

write $\phi_{,x} := \frac{\partial \phi}{\partial x} =: (\tilde{d}\phi)_x$

ϕ depends either on x and y , or on x' and y'

$$\Rightarrow \phi_{,x'} = \phi_{,x} x_{,x'} + \phi_{,y} y_{,x'}$$

$$(\phi_{,x'}, \phi_{,y'}) = (\phi_{,x} x_{,x'} + \phi_{,y} y_{,x'}, \phi_{,x} x_{,y'} + \phi_{,y} y_{,y'})$$

GR: coord. transf.: 1-forms

ϕ = scalar field = $\phi(x, y) \equiv \phi(x', y')$

write $\phi_{,x} := \frac{\partial \phi}{\partial x} =: (\tilde{d}\phi)_x$

ϕ depends either on x and y , or on x' and y'

$$\Rightarrow \phi_{,x'} = \phi_{,x} x_{,x'} + \phi_{,y} y_{,x'}$$

$$(\phi_{,x'}, \phi_{,y'}) = (\phi_{,x}, \phi_{,y}) \begin{pmatrix} x_{,x'} & x_{,y'} \\ y_{,x'} & y_{,y'} \end{pmatrix}$$

GR: coord. transf.: 1-forms

ϕ = scalar field = $\phi(x, y) \equiv \phi(x', y')$

write $\phi_{,x} := \frac{\partial \phi}{\partial x} =: (\tilde{d}\phi)_x$

ϕ depends either on x and y , or on x' and y'

$$\Rightarrow \phi_{,x'} = \phi_{,x} x_{,x'} + \phi_{,y} y_{,x'}$$

$$(\phi_{,x'}, \phi_{,y'}) = (\phi_{,x}, \phi_{,y}) \begin{pmatrix} x_{,x'} & x_{,y'} \\ y_{,x'} & y_{,y'} \end{pmatrix}$$

$$\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} x' \\ y' \end{pmatrix} \quad (\text{example: rotation})$$

$$x_{,x'} = \frac{\partial x}{\partial x'} = \cos \theta$$

$$x_{,y'} = \frac{\partial x}{\partial y'} = -\sin \theta \dots$$

GR: coord. transf.: 1-forms

ϕ = scalar field = $\phi(x, y) \equiv \phi(x', y')$

write $\phi_{,x} := \frac{\partial \phi}{\partial x} =: (\tilde{d}\phi)_x$

ϕ depends either on x and y , or on x' and y'

$$\Rightarrow \phi_{,x'} = \phi_{,x} x_{,x'} + \phi_{,y} y_{,x'}$$

$$(\phi_{,x'}, \phi_{,y'}) = (\phi_{,x}, \phi_{,y}) \begin{pmatrix} x_{,x'} & x_{,y'} \\ y_{,x'} & y_{,y'} \end{pmatrix}$$

$$\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x_{,x'} & x_{,y'} \\ y_{,x'} & y_{,y'} \end{pmatrix} \begin{pmatrix} x' \\ y' \end{pmatrix} \quad (\text{general})$$

GR: coord. transf.: 1-forms

ϕ = scalar field = $\phi(x, y) \equiv \phi(x', y')$

write $\phi_{,x} := \frac{\partial \phi}{\partial x} =: (\tilde{d}\phi)_x$

ϕ depends either on x and y , or on x' and y'

$$\Rightarrow \phi_{,x'} = \phi_{,x} x_{,x'} + \phi_{,y} y_{,x'}$$

$$(\phi_{,x'}, \phi_{,y'}) = (\phi_{,x}, \phi_{,y}) \begin{pmatrix} x_{,x'} & x_{,y'} \\ y_{,x'} & y_{,y'} \end{pmatrix}$$

$$\begin{pmatrix} x \\ y \end{pmatrix} = \Lambda^{-1} \begin{pmatrix} x' \\ y' \end{pmatrix} \quad (\text{general})$$

GR: coord. transf.: 1-forms

ϕ = scalar field = $\phi(x, y) \equiv \phi(x', y')$

write $\phi_{,x} := \frac{\partial \phi}{\partial x} =: (\tilde{d}\phi)_x$

ϕ depends either on x and y , or on x' and y'

$$\Rightarrow \phi_{,x'} = \phi_{,x} x_{,x'} + \phi_{,y} y_{,x'}$$

$$(\phi_{,x'}, \phi_{,y'}) = (\phi_{,x}, \phi_{,y}) \begin{pmatrix} x_{,x'} & x_{,y'} \\ y_{,x'} & y_{,y'} \end{pmatrix}$$

$$\begin{pmatrix} x \\ y \end{pmatrix} = \Lambda^{-1} \begin{pmatrix} x' \\ y' \end{pmatrix} \quad (\text{general})$$

$$\Rightarrow (\phi_{,x'}, \phi_{,y'}) = (\phi_{,x}, \phi_{,y}) \Lambda^{-1}$$

GR: coord. transf.: 1-forms

ϕ = scalar field = $\phi(x, y) \equiv \phi(x', y')$

write $\phi_{,x} := \frac{\partial \phi}{\partial x} =: (\tilde{d}\phi)_x$

ϕ depends either on x and y , or on x' and y'

$$\Rightarrow \phi_{,x'} = \phi_{,x} x_{,x'} + \phi_{,y} y_{,x'}$$

$$(\phi_{,x'}, \phi_{,y'}) = (\phi_{,x}, \phi_{,y}) \begin{pmatrix} x_{,x'} & x_{,y'} \\ y_{,x'} & y_{,y'} \end{pmatrix}$$

$$\begin{pmatrix} x \\ y \end{pmatrix} = \Lambda^{-1} \begin{pmatrix} x' \\ y' \end{pmatrix} \quad (\text{general})$$

$$\tilde{d}\phi = \left((\tilde{d}\phi)_{x'}, (\tilde{d}\phi)_{y'} \right) = \left((\tilde{d}\phi)_x, (\tilde{d}\phi)_y \right) \Lambda^{-1}$$

GR: coord. transf.: 1-forms

ϕ = scalar field = $\phi(x, y) \equiv \phi(x', y')$

write $\phi_{,x} := \frac{\partial \phi}{\partial x} =: (\tilde{d}\phi)_x$

ϕ depends either on x and y , or on x' and y'

$$\Rightarrow \phi_{,x'} = \phi_{,x} x_{,x'} + \phi_{,y} y_{,x'}$$

$$(\phi_{,x'}, \phi_{,y'}) = (\phi_{,x}, \phi_{,y}) \begin{pmatrix} x_{,x'} & x_{,y'} \\ y_{,x'} & y_{,y'} \end{pmatrix}$$

$$\begin{pmatrix} x \\ y \end{pmatrix} = \Lambda^{-1} \begin{pmatrix} x' \\ y' \end{pmatrix} \quad (\text{general})$$

$$\tilde{d}\phi = ((\tilde{d}\phi)_{x'}, (\tilde{d}\phi)_{y'}) = ((\tilde{d}\phi)_x, (\tilde{d}\phi)_y) \Lambda^{-1}$$

$$(\tilde{d}\phi)_{\mu'} = (\tilde{d}\phi)_{\nu} \Lambda^{\nu}_{\mu'}$$

GR: coord. transf.: 1-forms

basis vectors of different bases: $\vec{e}_{\mu'} = \Lambda^{\nu}_{\mu'} \vec{e}_{\nu}$

same vector: $(\vec{p})^{\mu'} = \Lambda^{\mu'}_{\nu} (\vec{p})^{\nu}$

GR: coord. transf.: 1-forms

basis vectors of different bases: $\vec{e}_{\mu'} = \Lambda^{\nu}_{\mu'} \vec{e}_{\nu}$

same vector: $p^{\mu'} = \Lambda^{\mu'}_{\nu} p^{\nu}$

same gradient (example 1-form): $(\tilde{d}\phi)_{\mu'} = (\tilde{d}\phi)_{\nu} \Lambda^{\nu}_{\mu'}$

GR: coord. transf.: 1-forms

basis vectors of different bases: $\vec{e}_{\mu'} = \Lambda_{\mu'}^{\nu} \vec{e}_{\nu}$

same vector: $p^{\mu'} = \Lambda_{\nu}^{\mu'} p^{\nu}$

same gradient (example 1-form): $(\tilde{d}\phi)_{\mu'} = (\tilde{d}\phi)_{\nu} \Lambda_{\mu'}^{\nu}$

- vector $\vec{p}$ is **in**variant: no dependence on coords
- $\vec{p}$ is **contra**variant: components p^{ν} change inversely to how $\vec{e}_{\mu}$ change; inverses: matrix $\{\Lambda_{\mu'}^{\nu}\}$ vs $\{\Lambda_{\alpha}^{\beta'}\}$

GR: coord. transf.: 1-forms

basis vectors of different bases: $\vec{e}_{\mu'} = \Lambda^{\nu}_{\mu'} \vec{e}_{\nu}$

same vector: $p^{\mu'} = \Lambda^{\mu'}_{\nu} p^{\nu}$

same gradient (example 1-form): $(\tilde{d}\phi)_{\mu'} = (\tilde{d}\phi)_{\nu} \Lambda^{\nu}_{\mu'}$

- vector $\vec{p}$ is **in**variant: no dependence on coords
- $\vec{p}$ is **contra**variant: components p^{ν} change inversely to how $\vec{e}_{\mu}$ change; inverses: matrix $\{\Lambda^{\nu}_{\mu'}\}$ vs $\{\Lambda^{\beta'}_{\alpha}\}$
- 1-form $\tilde{d}\phi$ is **in**variant: no dependence on coords
- $\tilde{d}\phi$ is **cova**riant: components $(\tilde{d}\phi)_{\mu}$ change like $\vec{e}_{\mu}$ (but left-multiply)

GR: coord. transf.: 1-forms

basis vectors of different bases: $\vec{e}_{\mu'} = \Lambda^{\nu}_{\mu'} \vec{e}_{\nu}$

same vector: $p^{\mu'} = \Lambda^{\mu'}_{\nu} p^{\nu}$

same gradient (example 1-form): $(\tilde{d}\phi)_{\mu'} = (\tilde{d}\phi)_{\nu} \Lambda^{\nu}_{\mu'}$

- vector $\vec{p}$ is **in**variant: no dependence on coords
- $\vec{p}$ is **contra**variant: components p^{ν} change inversely to

how $\vec{e}_{\mu}$ change; inverses: matrix $\{\Lambda^{\nu}_{\mu'}\}$ vs $\{\Lambda^{\beta'}_{\alpha}\}$

- 1-form $\tilde{d}\phi$ is **in**variant: no dependence on coords
- $\tilde{d}\phi$ is **cova**riant: components $(\tilde{d}\phi)_{\mu}$ change like $\vec{e}_{\mu}$ (but left-multiply)

w:Covariance and contravariance of vectors


$$\mathbf{GR}: \vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, \mathbf{g}$$

GR tensors: two different scalar products





$$\mathbf{GR}: \vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, \mathbf{g}$$

GR tensors: two different scalar products
vector–1-form duality requirement:





$$\mathbf{GR}: \vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, g$$

GR tensors: two different scalar products

vector–1-form duality requirement:

$$\langle \vec{p}, \tilde{q} \rangle = \sum_{\mu} p^{\mu} q_{\mu}$$





$$\mathbf{GR}: \vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, g$$

GR tensors: two different scalar products

vector–1-form duality requirement:

$$\langle \vec{p}, \tilde{q} \rangle = p^\mu q_\mu$$





$$\mathbf{GR}: \vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, g$$

GR tensors: two different scalar products

vector–1-form duality requirement:

$$\langle \vec{p}, \tilde{q} \rangle = p^\mu q_\mu = \vec{p}(\tilde{q})$$





$$\mathbf{GR}: \vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, g$$

GR tensors: two different scalar products

vector–1-form duality requirement:

$$\langle \vec{p}, \tilde{q} \rangle = p^\mu q_\mu = \vec{p}(\tilde{q}) = \tilde{q}(\vec{p})$$





GR: $\vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, g$

GR tensors: two different scalar products

vector–1-form duality requirement:

$$\langle \vec{p}, \tilde{q} \rangle = p^\mu q_\mu = \vec{p}(\tilde{q}) = \tilde{q}(\vec{p})$$

$\langle , \rangle$ is a (1,1) tensor



GR: $\vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, g$

GR tensors: two different scalar products

vector–1-form duality requirement:

$$\langle \vec{p}, \tilde{q} \rangle = p^\mu q_\mu = \vec{p}(\tilde{q}) = \tilde{q}(\vec{p})$$

$\langle , \rangle$ is a (1,1) tensor

can be called I with components δ^μ_ν in a coordinate basis

GR: $\vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, g$

GR tensors: two different scalar products

vector–1-form duality requirement:

$$\langle \vec{p}, \tilde{q} \rangle = p^\mu q_\mu = \vec{p}(\tilde{q}) = \tilde{q}(\vec{p})$$

$\langle , \rangle$ is a (1,1) tensor

think: vector $\rightarrow$ column vector

1-form $\rightarrow$ row vector



$$\mathbf{GR}: \vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, \mathbf{g}$$

GR tensors: two different scalar products

vector–1-form duality requirement:

$$\langle \vec{p}, \tilde{q} \rangle = p^\mu q_\mu = \vec{p}(\tilde{q}) = \tilde{q}(\vec{p})$$

$\langle , \rangle$ is a (1,1) tensor

$$(q_0, q_1) \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} p^0 \\ p^1 \end{pmatrix} =$$



GR: $\vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, g$

GR tensors: two different scalar products

vector–1-form duality requirement:

$$\langle \vec{p}, \tilde{q} \rangle = p^\mu q_\mu = \vec{p}(\tilde{q}) = \tilde{q}(\vec{p})$$

$\langle , \rangle$ is a (1,1) tensor

$$(q_0, q_1) \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} p^0 \\ p^1 \end{pmatrix} = (q_0, q_1) \begin{pmatrix} p^0 \\ p^1 \end{pmatrix}$$

GR: $\vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, g$

GR tensors: two different scalar products

vector–1-form duality requirement:

$$\langle \vec{p}, \tilde{q} \rangle = p^\mu q_\mu = \vec{p}(\tilde{q}) = \tilde{q}(\vec{p})$$

$\langle , \rangle$ is a (1,1) tensor

$$(q_0, q_1) \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} p^0 \\ p^1 \end{pmatrix} = (q_0, q_1) \begin{pmatrix} p^0 \\ p^1 \end{pmatrix} = p^\mu q_\mu$$

GR: $\vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, g$

GR tensors: two different scalar products

vector–1-form duality requirement:

$$\langle \vec{p}, \tilde{q} \rangle = p^\mu q_\mu = \vec{p}(\tilde{q}) = \tilde{q}(\vec{p})$$

$\langle , \rangle$ is a (1,1) tensor

$$(q_0, q_1) \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} p^0 \\ p^1 \end{pmatrix} = (q_0, q_1) \begin{pmatrix} p^0 \\ p^1 \end{pmatrix} = p^\mu q_\mu$$

$\langle , \rangle = (1,1)$ -tensor = “row-column” matrix I with $I^\mu_\nu = \delta^\mu_\nu$



GR: $\vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, \mathbf{g}$

GR tensors: two different scalar products





GR: $\vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, g$

GR tensors: two different scalar products

ordinary linear algebra: column vectors, row vectors,
matrices





GR: $\vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, \mathbf{g}$

GR tensors: two different scalar products

(m, n) -tensor algebra: m column n row $m + n$ -arrays





GR: $\vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, g$

GR tensors: two different scalar products

(m, n) -tensor algebra: m column n row $m + n$ -arrays

e.g.: $(0, 2)$ -tensor: metric $g_{\mu\nu}$





GR: $\vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, \mathbf{g}$

GR tensors: two different scalar products

(m, n) -tensor algebra: m column n row $m + n$ -arrays

using $\langle, \rangle$, $(1, 0)$ -tensor = vector = function of 1-forms





GR: $\vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, \mathbf{g}$

GR tensors: two different scalar products

(m, n) -tensor algebra: m column n row $m + n$ -arrays

using $\langle, \rangle$, $(0, 1)$ -tensor = 1-form = function of vectors



GR: $\vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, g$

GR tensors: two different scalar products

(m, n) -tensor algebra: m column n row $m + n$ -arrays

(m, n) -tensor = function of m 1-forms and n vectors

GR: $\vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, g$

GR tensors: two different scalar products

(m, n) -tensor algebra: m column n row $m + n$ -arrays

(m, n) -tensor = function of m 1-forms and n vectors

V = space of vectors $\vec{p} = p^\mu \vec{e}_\mu$

GR: $\vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, g$

GR tensors: two different scalar products

(m, n) -tensor algebra: m column n row $m + n$ -arrays

(m, n) -tensor = function of m 1-forms and n vectors

V = space of vectors $\vec{p} = p^\mu \vec{e}_\mu$

V^* = dual space of 1-forms $\tilde{q} = q_\mu \tilde{e}^\mu$

GR: $\vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, g$

GR tensors: two different scalar products

(m, n) -tensor algebra: m column n row $m + n$ -arrays

(m, n) -tensor = function of m 1-forms and n vectors

V = space of vectors $\vec{p} = p^\mu \vec{e}_\mu$

V^* = dual space of 1-forms $\tilde{q} = q_\mu \tilde{e}^\mu$

$V^* \otimes V^*$ = space of $(0, 2)$ -tensors $T = T_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$ (e.g. metric) w:tensor product

GR: $\vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, g$

GR tensors: two different scalar products

(m, n) -tensor algebra: m column n row $m + n$ -arrays

(m, n) -tensor = function of m 1-forms and n vectors

V = space of vectors $\vec{p} = p^\mu \vec{e}_\mu$

V^* = dual space of 1-forms $\tilde{q} = q_\mu \tilde{e}^\mu$

$V^* \otimes V^*$ = space of $(0, 2)$ -tensors $T = T_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$ (e.g. metric) w:tensor product

loosely speaking, the second $\otimes$ means “function of two vectors” (or 1-forms, or a vector and a 1-form) in *that particular left-to-right order*

GR: $\vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, g$

GR tensors: two different scalar products

(m, n) -tensor algebra: m column n row $m + n$ -arrays

(m, n) -tensor = function of m 1-forms and n vectors

V = space of vectors $\vec{p} = p^\mu \vec{e}_\mu$

V^* = dual space of 1-forms $\tilde{q} = q_\mu \tilde{e}^\mu$

$V^* \otimes V^*$ = space of $(0, 2)$ -tensors $T = T_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$ (e.g. metric) w:tensor product

order of $V^* \otimes V^* = 2$

GR: $\vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, g$

GR tensors: two different scalar products

(m, n) -tensor algebra: m column n row $m + n$ -arrays

(m, n) -tensor = function of m 1-forms and n vectors

V = space of vectors $\vec{p} = p^\mu \vec{e}_\mu$

V^* = dual space of 1-forms $\tilde{q} = q_\mu \tilde{e}^\mu$

$V^* \otimes V^*$ = space of $(0, 2)$ -tensors $T = T_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$ (e.g. metric) w:tensor product

order of $V^* \otimes V^* = 2$

warning: the "rank" of tensors has two different meanings: w:Tensor_(intrinsic_definition)#Tensor_rank

GR: $\vec{p}, \tilde{q}, \langle \vec{p}, \tilde{q} \rangle, g$

GR tensors: two different scalar products

(m, n) -tensor algebra: m column n row $m + n$ -arrays

(m, n) -tensor = function of m 1-forms and n vectors

V = space of vectors $\vec{p} = p^\mu \vec{e}_\mu$

V^* = dual space of 1-forms $\tilde{q} = q_\mu \tilde{e}^\mu$

$V^* \otimes V^*$ = space of $(0, 2)$ -tensors $T = T_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$ (e.g. metric) w:tensor product

order of $V^* \otimes V^* = 2$

dimension of $V^* \otimes V^* = 16$ (for V = spacetime)

GR: g

$V^* \otimes V^*$ = space of $(0, 2)$ -tensors $T = T_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$, where
 $\otimes =$ w:tensor product

GR: g

$V^* \otimes V^*$ = space of $(0, 2)$ -tensors $T = T_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$, where

$\otimes =$ w:tensor product

e.g.: metric g = function of two vectors

GR: g

$V^* \otimes V^*$ = space of $(0, 2)$ -tensors $T = T_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$, where

$\otimes =$ w:tensor product

e.g.: metric g = function of two vectors

= “row-row” matrix

GR: g

$V^* \otimes V^*$ = space of $(0, 2)$ -tensors $T = T_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$, where
 $\otimes =$ w:tensor product

e.g.: metric g = function of two vectors

= “row-row” matrix

e.g. Euclidean g on $\mathbb{R}^2$. $g_{r\theta} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}$

GR: g

$V^* \otimes V^*$ = space of $(0, 2)$ -tensors $T = T_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$, where
 $\otimes =$ w:tensor product

e.g.: metric g = function of two vectors
= “row-row” matrix

e.g. Euclidean g on $\mathbb{R}^2$. $g_{r\theta} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}$ or

$$g_{xy} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

GR: g

$V^* \otimes V^*$ = space of $(0, 2)$ -tensors $T = T_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$, where
 $\otimes =$ w:tensor product

e.g.: metric g = function of two vectors

= “row-row” matrix

e.g. Euclidean g on $\mathbb{R}^2$. $g_{r\theta} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}$ or

$$g_{xy} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$g(\vec{A}, \vec{B})$$

GR: g

$V^* \otimes V^*$ = space of $(0, 2)$ -tensors $T = T_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$, where
 $\otimes =$ w:tensor product

e.g.: metric g = function of two vectors
= “row-row” matrix

e.g. Euclidean g on $\mathbb{R}^2$. $g_{r\theta} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}$ or

$$g_{xy} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

also written: $\vec{A} \cdot \vec{B}$ “dot product”

GR: g

$V^* \otimes V^*$ = space of $(0, 2)$ -tensors $T = T_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$, where
 $\otimes =$ w:tensor product

e.g.: metric g = function of two vectors

= “row-row” matrix

e.g. Euclidean g on $\mathbb{R}^2$. $g_{r\theta} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}$ or

$$g_{xy} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$g(\vec{A}, \vec{B})$$

GR: g

$V^* \otimes V^*$ = space of $(0, 2)$ -tensors $T = T_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$, where
 $\otimes =$ w:tensor product

e.g.: metric g = function of two vectors
= “row-row” matrix

e.g. Euclidean g on $\mathbb{R}^2$. $g_{r\theta} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}$ or

$$g_{xy} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\mathbf{g}(\vec{A}, \vec{B}) = \left[\begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix} \begin{pmatrix} A_r \\ A_\theta \end{pmatrix} \right]^T \begin{pmatrix} B_r \\ B_\theta \end{pmatrix}$$

GR: g

$V^* \otimes V^*$ = space of $(0, 2)$ -tensors $T = T_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$, where
 $\otimes =$ w:tensor product

e.g.: metric g = function of two vectors
= “row-row” matrix

e.g. Euclidean g on $\mathbb{R}^2$. $g_{r\theta} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}$ or

$$g_{xy} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\mathbf{g}(\vec{A}, \vec{B}) = (A_r, A_\theta r^2) \begin{pmatrix} B_r \\ B_\theta \end{pmatrix}$$

GR: g

$V^* \otimes V^*$ = space of $(0, 2)$ -tensors $T = T_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$, where
 $\otimes =$ w:tensor product

e.g.: metric g = function of two vectors
= “row-row” matrix

e.g. Euclidean g on $\mathbb{R}^2$. $g_{r\theta} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}$ or

$$g_{xy} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\mathbf{g}(\vec{A}, \vec{B}) = A_r B_r + A_\theta B_\theta r^2$$

GR: g

$V^* \otimes V^*$ = space of $(0, 2)$ -tensors $T = T_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$, where
 $\otimes =$ w:tensor product

e.g.: metric g = function of two vectors
= “row-row” matrix

e.g. Euclidean g on $\mathbb{R}^2$. $g_{r\theta} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}$ or

$$g_{xy} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\mathbf{g}(\vec{A}, \vec{B}) = A_r B_r + A_\theta B_\theta r^2$$

$$= \left[\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} A_x \\ A_y \end{pmatrix} \right]^T \begin{pmatrix} B_x \\ B_y \end{pmatrix}$$

GR: g

$V^* \otimes V^*$ = space of $(0, 2)$ -tensors $T = T_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$, where
 $\otimes =$ w:tensor product

e.g.: metric g = function of two vectors
= “row-row” matrix

e.g. Euclidean g on $\mathbb{R}^2$. $g_{r\theta} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}$ or

$$g_{xy} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\mathbf{g}(\vec{A}, \vec{B}) = A_r B_r + A_\theta B_\theta r^2 = (A_x, A_y) \begin{pmatrix} B_x \\ B_y \end{pmatrix}$$

GR: g

$V^* \otimes V^*$ = space of $(0, 2)$ -tensors $T = T_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$, where
 $\otimes =$ w:tensor product

e.g.: metric g = function of two vectors
= “row-row” matrix

e.g. Euclidean g on $\mathbb{R}^2$. $g_{r\theta} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}$ or

$$g_{xy} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\mathbf{g}(\vec{A}, \vec{B}) = A_r B_r + A_\theta B_\theta r^2 = A_x B_x + A_y B_y$$

GR: g

$V^* \otimes V^*$ = space of $(0, 2)$ -tensors $T = T_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$, where
 $\otimes =$ w:tensor product

e.g.: metric g = function of two vectors

= “row-row” matrix

e.g. Euclidean g on $\mathbb{R}^2$. $g_{r\theta} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}$ or

$$g_{xy} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\mathbf{g}(\vec{A}, \vec{B}) = A_r B_r + A_\theta B_\theta r^2 = A_x B_x + A_y B_y$$

in general, for a 2-form T , $T(\vec{A}, \vec{B}) \neq T(\vec{B}, \vec{A})$

GR: g

$V^* \otimes V^*$ = space of $(0, 2)$ -tensors $T = T_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$, where
 $\otimes =$ w:tensor product

e.g.: metric g = function of two vectors
= “row-row” matrix

e.g. Euclidean g on $\mathbb{R}^2$. $g_{r\theta} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}$ or

$$g_{xy} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\mathbf{g}(\vec{A}, \vec{B}) = A_r B_r + A_\theta B_\theta r^2 = A_x B_x + A_y B_y$$

$$\mathbf{g} = g_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$$

GR: g

$V^* \otimes V^*$ = space of $(0, 2)$ -tensors $T = T_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$, where
 $\otimes =$ w:tensor product

e.g.: metric g = function of two vectors
= “row-row” matrix

e.g. Euclidean g on $\mathbb{R}^2$. $g_{r\theta} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}$ or

$$g_{xy} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\mathbf{g}(\vec{A}, \vec{B}) = A_r B_r + A_\theta B_\theta r^2 = A_x B_x + A_y B_y$$

$$\mathbf{g} = g_{r\theta} \tilde{e}^r \otimes \tilde{e}^\theta$$

GR: g

$V^* \otimes V^*$ = space of $(0, 2)$ -tensors $T = T_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$, where
 $\otimes =$ w:tensor product

e.g.: metric g = function of two vectors
= “row-row” matrix

e.g. Euclidean g on $\mathbb{R}^2$. $g_{r\theta} = \begin{pmatrix} 1 & 0 \\ 0 & r^2 \end{pmatrix}$ or

$$g_{xy} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$\mathbf{g}(\vec{A}, \vec{B}) = A_r B_r + A_\theta B_\theta r^2 = A_x B_x + A_y B_y$$

$$\mathbf{g} = g_{xy} \tilde{e}^x \otimes \tilde{e}^y$$

GR: metric tensor g , g^{-1} , bases

g can be applied to basis vectors $\vec{e}_\mu$

GR: metric tensor g , g^{-1} , bases

g can be applied to basis vectors $\vec{e}_\mu$

we can define components (used earlier): $g_{\mu\nu} := g(\vec{e}_\mu, \vec{e}_\nu)$

GR: metric tensor g , g^{-1} , bases

g can be applied to basis vectors $\vec{e}_\mu$

we can define components (used earlier): $g_{\mu\nu} := g(\vec{e}_\mu, \vec{e}_\nu)$

$$\Rightarrow g = g_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$$

GR: metric tensor g , g^{-1} , bases

g can be applied to basis vectors $\vec{e}_\mu$

we can define components (used earlier): $g_{\mu\nu} := g(\vec{e}_\mu, \vec{e}_\nu)$

$$\Rightarrow g = g_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$$

$$\text{e.g. } g = g_{rr} \tilde{e}^r \otimes \tilde{e}^r + g_{r\theta} \tilde{e}^r \otimes \tilde{e}^\theta + g_{\theta r} \tilde{e}^\theta \otimes \tilde{e}^r + g_{\theta\theta} \tilde{e}^\theta \otimes \tilde{e}^\theta$$

GR: metric tensor g , g^{-1} , bases

g can be applied to basis vectors $\vec{e}_\mu$

we can define components (used earlier): $g_{\mu\nu} := g(\vec{e}_\mu, \vec{e}_\nu)$

$$\Rightarrow g = g_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$$

$$\text{e.g. } g = g_{rr} \tilde{e}^r \otimes \tilde{e}^r + g_{\theta\theta} \tilde{e}^\theta \otimes \tilde{e}^\theta$$

GR: metric tensor g , g^{-1} , bases

g can be applied to basis vectors $\vec{e}_\mu$

we can define components (used earlier): $g_{\mu\nu} := g(\vec{e}_\mu, \vec{e}_\nu)$

$$\Rightarrow g = g_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$$

$$\text{e.g. } g = g_{rr} \tilde{e}^r \otimes \tilde{e}^r + g_{\theta\theta} \tilde{e}^\theta \otimes \tilde{e}^\theta$$

check: $g(\vec{e}_r, \vec{e}_r) = g_{rr}$?

GR: metric tensor g , g^{-1} , bases

g can be applied to basis vectors $\vec{e}_\mu$

we can define components (used earlier): $g_{\mu\nu} := g(\vec{e}_\mu, \vec{e}_\nu)$

$$\Rightarrow g = g_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$$

$$\text{e.g. } g = g_{rr} \tilde{e}^r \otimes \tilde{e}^r + g_{\theta\theta} \tilde{e}^\theta \otimes \tilde{e}^\theta$$

$$g(\vec{e}_r, \vec{e}_r) = (g_{rr} \tilde{e}^r \otimes \tilde{e}^r + g_{\theta\theta} \tilde{e}^\theta \otimes \tilde{e}^\theta)(\vec{e}_r, \vec{e}_r)$$

GR: metric tensor g , g^{-1} , bases

g can be applied to basis vectors $\vec{e}_\mu$

we can define components (used earlier): $g_{\mu\nu} := g(\vec{e}_\mu, \vec{e}_\nu)$

$$\Rightarrow g = g_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$$

$$\text{e.g. } g = g_{rr} \tilde{e}^r \otimes \tilde{e}^r + g_{\theta\theta} \tilde{e}^\theta \otimes \tilde{e}^\theta$$

$$g(\vec{e}_r, \vec{e}_r) = g_{rr} \tilde{e}^r \otimes \tilde{e}^r(\vec{e}_r, \vec{e}_r) + g_{\theta\theta} \tilde{e}^\theta \otimes \tilde{e}^\theta(\vec{e}_r, \vec{e}_r)$$

GR: metric tensor g , g^{-1} , bases

g can be applied to basis vectors $\vec{e}_\mu$

we can define components (used earlier): $g_{\mu\nu} := g(\vec{e}_\mu, \vec{e}_\nu)$

$$\Rightarrow g = g_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$$

$$\text{e.g. } g = g_{rr} \tilde{e}^r \otimes \tilde{e}^r + g_{\theta\theta} \tilde{e}^\theta \otimes \tilde{e}^\theta$$

$$g(\vec{e}_r, \vec{e}_r) = g_{rr} \tilde{e}^r(\vec{e}_r) \tilde{e}^r(\vec{e}_r) + g_{\theta\theta} \tilde{e}^\theta(\vec{e}_r) \tilde{e}^\theta(\vec{e}_r)$$

GR: metric tensor g , g^{-1} , bases

g can be applied to basis vectors $\vec{e}_\mu$

we can define components (used earlier): $g_{\mu\nu} := g(\vec{e}_\mu, \vec{e}_\nu)$

$$\Rightarrow g = g_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$$

$$\text{e.g. } g = g_{rr} \tilde{e}^r \otimes \tilde{e}^r + g_{\theta\theta} \tilde{e}^\theta \otimes \tilde{e}^\theta$$

$$g(\vec{e}_r, \vec{e}_r) = g_{rr} \langle \tilde{e}^r, \vec{e}_r \rangle \langle \tilde{e}^r, \vec{e}_r \rangle + g_{\theta\theta} \langle \tilde{e}^\theta, \vec{e}_r \rangle \langle \tilde{e}^\theta, \vec{e}_r \rangle$$

GR: metric tensor g , g^{-1} , bases

g can be applied to basis vectors $\vec{e}_\mu$

we can define components (used earlier): $g_{\mu\nu} := g(\vec{e}_\mu, \vec{e}_\nu)$

$$\Rightarrow g = g_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$$

$$\text{e.g. } g = g_{rr} \tilde{e}^r \otimes \tilde{e}^r + g_{\theta\theta} \tilde{e}^\theta \otimes \tilde{e}^\theta$$

$g(\vec{e}_r, \vec{e}_r) = g_{rr} \times 1 \times 1 + g_{\theta\theta} \times 0 \times 0$ by duality through scalar product $\langle , \rangle$

GR: metric tensor g , g^{-1} , bases

g can be applied to basis vectors $\vec{e}_\mu$

we can define components (used earlier): $g_{\mu\nu} := g(\vec{e}_\mu, \vec{e}_\nu)$

$$\Rightarrow g = g_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$$

$$\text{e.g. } g = g_{rr} \tilde{e}^r \otimes \tilde{e}^r + g_{\theta\theta} \tilde{e}^\theta \otimes \tilde{e}^\theta$$

$$g(\vec{e}_r, \vec{e}_r) = g_{rr} \text{ self-consistent definition}$$

GR: metric tensor g , g^{-1} , bases

g can be applied to basis vectors $\vec{e}_\mu$

we can define components (used earlier): $g_{\mu\nu} := g(\vec{e}_\mu, \vec{e}_\nu)$

$$\Rightarrow g = g_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$$

$$\text{e.g. } g = g_{rr} \tilde{e}^r \otimes \tilde{e}^r + g_{\theta\theta} \tilde{e}^\theta \otimes \tilde{e}^\theta$$

$$\text{inverse: } g^{-1} = g^{\mu\nu} \vec{e}_\mu \otimes \vec{e}_\nu,$$

GR: metric tensor g , g^{-1} , bases

g can be applied to basis vectors $\vec{e}_\mu$

we can define components (used earlier): $g_{\mu\nu} := g(\vec{e}_\mu, \vec{e}_\nu)$

$$\Rightarrow g = g_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$$

$$\text{e.g. } g = g_{rr} \tilde{e}^r \otimes \tilde{e}^r + g_{\theta\theta} \tilde{e}^\theta \otimes \tilde{e}^\theta$$

$$\text{inverse: } g^{-1} = g^{\mu\nu} \vec{e}_\mu \otimes \vec{e}_\nu,$$

$$\text{where } g^{\mu\alpha} g_{\alpha\nu} = \delta^\mu_\nu$$

GR: metric tensor g , g^{-1} , bases

g can be applied to basis vectors $\vec{e}_\mu$

we can define components (used earlier): $g_{\mu\nu} := g(\vec{e}_\mu, \vec{e}_\nu)$

$$\Rightarrow g = g_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$$

$$\text{e.g. } g = g_{rr} \tilde{e}^r \otimes \tilde{e}^r + g_{\theta\theta} \tilde{e}^\theta \otimes \tilde{e}^\theta$$

$$\text{inverse: } g^{-1} = g^{\mu\nu} \vec{e}_\mu \otimes \vec{e}_\nu,$$

$$\text{where } g^{\mu\alpha} g_{\alpha\nu} = \delta^\mu_\nu$$

duality of associate vectors and 1-forms:

$$g(\vec{A}, \vec{B}) = g^{-1}(\tilde{A}, \tilde{B})$$

GR: metric tensor g , g^{-1} , bases

g can be applied to basis vectors $\vec{e}_\mu$

we can define components (used earlier): $g_{\mu\nu} := g(\vec{e}_\mu, \vec{e}_\nu)$

$$\Rightarrow g = g_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$$

$$\text{e.g. } g = g_{rr} \tilde{e}^r \otimes \tilde{e}^r + g_{\theta\theta} \tilde{e}^\theta \otimes \tilde{e}^\theta$$

$$\text{inverse: } g^{-1} = g^{\mu\nu} \vec{e}_\mu \otimes \vec{e}_\nu,$$

$$\text{where } g^{\mu\alpha} g_{\alpha\nu} = \delta^\mu_\nu$$

duality of associate vectors and 1-forms:

$$g(\vec{A}, \vec{B}) = g^{-1}(\tilde{A}, \tilde{B}) = \vec{A} \cdot \vec{B}$$

GR: metric tensor g , g^{-1} , bases

g can be applied to basis vectors $\vec{e}_\mu$

we can define components (used earlier): $g_{\mu\nu} := g(\vec{e}_\mu, \vec{e}_\nu)$

$$\Rightarrow g = g_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$$

$$\text{e.g. } g = g_{rr} \tilde{e}^r \otimes \tilde{e}^r + g_{\theta\theta} \tilde{e}^\theta \otimes \tilde{e}^\theta$$

$$\text{inverse: } g^{-1} = g^{\mu\nu} \vec{e}_\mu \otimes \vec{e}_\nu,$$

$$\text{where } g^{\mu\alpha} g_{\alpha\nu} = \delta^\mu_\nu$$

duality of associate vectors and 1-forms:

$$g(\vec{A}, \vec{B}) = g^{-1}(\tilde{A}, \tilde{B}) = \vec{A} \cdot \vec{B} = g_{\mu\nu} A^\mu B^\nu$$

GR: metric tensor g , g^{-1} , bases

g can be applied to basis vectors $\vec{e}_\mu$

we can define components (used earlier): $g_{\mu\nu} := g(\vec{e}_\mu, \vec{e}_\nu)$

$$\Rightarrow g = g_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$$

$$\text{e.g. } g = g_{rr} \tilde{e}^r \otimes \tilde{e}^r + g_{\theta\theta} \tilde{e}^\theta \otimes \tilde{e}^\theta$$

$$\text{inverse: } g^{-1} = g^{\mu\nu} \vec{e}_\mu \otimes \vec{e}_\nu,$$

$$\text{where } g^{\mu\alpha} g_{\alpha\nu} = \delta^\mu_\nu$$

duality of associate vectors and 1-forms:

$$g(\vec{A}, \vec{B}) = g^{-1}(\tilde{A}, \tilde{B}) = \vec{A} \cdot \vec{B} = g_{\mu\nu} A^\mu B^\nu = g^{\mu\nu} A_\mu B_\nu$$

GR: metric tensor g , g^{-1} , bases

g can be applied to basis vectors $\vec{e}_\mu$

we can define components (used earlier): $g_{\mu\nu} := g(\vec{e}_\mu, \vec{e}_\nu)$

$$\Rightarrow g = g_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$$

$$\text{e.g. } g = g_{rr} \tilde{e}^r \otimes \tilde{e}^r + g_{\theta\theta} \tilde{e}^\theta \otimes \tilde{e}^\theta$$

$$\text{inverse: } g^{-1} = g^{\mu\nu} \vec{e}_\mu \otimes \vec{e}_\nu,$$

$$\text{where } g^{\mu\alpha} g_{\alpha\nu} = \delta^\mu_\nu$$

duality of associate vectors and 1-forms:

$$g(\vec{A}, \vec{B}) = g^{-1}(\tilde{A}, \tilde{B}) = \vec{A} \cdot \vec{B} = g_{\mu\nu} A^\mu B^\nu = g^{\mu\nu} A_\mu B_\nu$$

$$\text{lower an index: } g_{\mu\nu} A^\mu = A_\nu$$

GR: metric tensor g , g^{-1} , bases

g can be applied to basis vectors $\vec{e}_\mu$

we can define components (used earlier): $g_{\mu\nu} := g(\vec{e}_\mu, \vec{e}_\nu)$

$$\Rightarrow g = g_{\mu\nu} \tilde{e}^\mu \otimes \tilde{e}^\nu$$

$$\text{e.g. } g = g_{rr} \tilde{e}^r \otimes \tilde{e}^r + g_{\theta\theta} \tilde{e}^\theta \otimes \tilde{e}^\theta$$

$$\text{inverse: } g^{-1} = g^{\mu\nu} \vec{e}_\mu \otimes \vec{e}_\nu,$$

$$\text{where } g^{\mu\alpha} g_{\alpha\nu} = \delta^\mu_\nu$$

duality of associate vectors and 1-forms:

$$g(\vec{A}, \vec{B}) = g^{-1}(\tilde{A}, \tilde{B}) = \vec{A} \cdot \vec{B} = g_{\mu\nu} A^\mu B^\nu = g^{\mu\nu} A_\mu B_\nu$$

$$\text{lower index: } g_{\mu\nu} A^\mu = A_\nu$$

$$\text{raise index: } g^{\mu\nu} B_\nu = B^\mu$$

GR: what is a coordinate?

a coordinate, e.g. x^0 or x^1 is a scalar field on the 4-manifold

GR: what is a coordinate?

a coordinate system x^μ = set of four scalar fields on the 4-manifold

GR: what is a coordinate?

a coordinate system x^μ = set of four scalar fields on the 4-manifold

(Bertschinger writes $x^\mu_{\mathbf{X}}$ to show dependence on position $\mathbf{X}$ in manifold $\neq$ vector space)

GR: what is a coordinate?


a coordinate system x^μ = set of four scalar fields on the 4-manifold

x^μ are differentiable *almost everywhere*

GR: what is a coordinate?

a coordinate system x^μ = set of four scalar fields on the 4-manifold

x^μ are differentiable *almost everywhere*




e.g. on $\mathbb{R}^2$

GR: what is a coordinate?

a coordinate system x^μ = set of four scalar fields on the 4-manifold

x^μ are differentiable *almost everywhere*




e.g. on $\mathbb{R}^2$

GR: what is a coordinate?

a coordinate system x^μ = set of four scalar fields on the 4-manifold

x^μ are differentiable *almost everywhere*




e.g. on $\mathbb{R}^2$

GR: what is a coordinate?

a coordinate system x^μ = set of four scalar fields on the 4-manifold

x^μ are differentiable *almost everywhere*




e.g. on $\mathbb{R}^2$

GR: what is a coordinate?

a coordinate system x^μ = set of four scalar fields on the 4-manifold

x^μ are differentiable *almost everywhere*



e.g. on $\mathbb{R}^2$

coordinate singularity $\neq$ singularity in manifold

GR: what is a coordinate basis?

coordinate basis: $\vec{e}_\mu, \tilde{e}^\nu$ chosen so that:

$$d\vec{x} = dx^\mu \vec{e}_\mu \text{ and}$$

GR: what is a coordinate basis?

coordinate basis: $\vec{e}_\mu, \tilde{e}^\nu$ chosen so that:

$$d\vec{x} = dx^\mu \vec{e}_\mu \text{ and}$$

$$df = \langle \tilde{d}f, d\vec{x} \rangle \text{ for any scalar field } f \text{ coordinate-free}$$

where $\tilde{d} = \tilde{e}^\mu \partial_\mu$ in a coordinate basis

GR: what is a coordinate basis?

coordinate basis: $\vec{e}_\mu, \tilde{e}^\nu$ chosen so that:

$d\vec{x} = dx^\mu \vec{e}_\mu$ and

$df = \langle \tilde{d}f, d\vec{x} \rangle$ for any scalar field f **coordinate-free**

where $\tilde{d} = \tilde{e}^\mu \partial_\mu$ in a coordinate basis

(Bertschinger writes $\tilde{\nabla}$ for the gradient $\tilde{d}$)

GR: what is a coordinate basis?

coordinate basis: $\vec{e}_\mu, \tilde{e}^\nu$ chosen so that:

$$d\vec{x} = dx^\mu \vec{e}_\mu \text{ and}$$

$$df = \langle \tilde{d}f, d\vec{x} \rangle \text{ for any scalar field } f \text{ coordinate-free}$$

where $\tilde{d} = \tilde{e}^\mu \partial_\mu$ in a coordinate basis

$$\text{check: } df = \langle \tilde{d}f, d\vec{x} \rangle$$

GR: what is a coordinate basis?

coordinate basis: $\vec{e}_\mu, \tilde{e}^\nu$ chosen so that:

$$d\vec{x} = dx^\mu \vec{e}_\mu \text{ and}$$

$$df = \langle \tilde{d}f, d\vec{x} \rangle \text{ for any scalar field } f \text{ coordinate-free}$$

where $\tilde{d} = \tilde{e}^\mu \partial_\mu$ in a coordinate basis

$$\begin{aligned} \text{check: } df &= \langle \tilde{d}f, d\vec{x} \rangle \\ &= \langle \tilde{e}^\mu \partial_\mu f, dx^\nu \vec{e}_\nu \rangle \end{aligned}$$

GR: what is a coordinate basis?

coordinate basis: $\vec{e}_\mu, \tilde{e}^\nu$ chosen so that:

$$d\vec{x} = dx^\mu \vec{e}_\mu \text{ and}$$

$$df = \langle \tilde{d}f, d\vec{x} \rangle \text{ for any scalar field } f \text{ coordinate-free}$$

where $\tilde{d} = \tilde{e}^\mu \partial_\mu$ in a coordinate basis

$$\text{check: } df = \langle \tilde{d}f, d\vec{x} \rangle$$

$$= \langle \tilde{e}^\mu \partial_\mu f, dx^\nu \vec{e}_\nu \rangle$$

$$= \partial_\mu f dx^\nu \langle \tilde{e}^\mu, \vec{e}_\nu \rangle \text{ since scalars commute}$$

GR: what is a coordinate basis?

coordinate basis: $\vec{e}_\mu, \tilde{e}^\nu$ chosen so that:

$$d\vec{x} = dx^\mu \vec{e}_\mu \text{ and}$$

$$df = \langle \tilde{d}f, d\vec{x} \rangle \text{ for any scalar field } f \text{ coordinate-free}$$

where $\tilde{d} = \tilde{e}^\mu \partial_\mu$ in a coordinate basis

$$\text{check: } df = \langle \tilde{d}f, d\vec{x} \rangle$$

$$= \langle \tilde{e}^\mu \partial_\mu f, dx^\nu \vec{e}_\nu \rangle$$

$$= \partial_\mu f dx^\nu \langle \tilde{e}^\mu, \vec{e}_\nu \rangle \text{ since scalars commute}$$

$$\text{i.e. } df = \partial_\mu f dx^\mu \text{ since } \langle \tilde{e}^\mu, \vec{e}_\nu \rangle = \delta_\nu^\mu$$

GR: what is a coordinate basis?

coordinate basis: $\vec{e}_\mu, \tilde{e}^\nu$ chosen so that:

$$d\vec{x} = dx^\mu \vec{e}_\mu \text{ and}$$

$$df = \langle \tilde{d}f, d\vec{x} \rangle \text{ for any scalar field } f \text{ coordinate-free}$$

where $\tilde{d} = \tilde{e}^\mu \partial_\mu$ in a coordinate basis

$$\text{check: } df = \langle \tilde{d}f, d\vec{x} \rangle$$

$$= \langle \tilde{e}^\mu \partial_\mu f, dx^\nu \vec{e}_\nu \rangle$$

$$= \partial_\mu f dx^\nu \langle \tilde{e}^\mu, \vec{e}_\nu \rangle \text{ since scalars commute}$$

$$\text{i.e. } df = \frac{\partial f}{\partial x^\mu} dx^\mu$$

GR: what is a coordinate basis?

coordinate basis: $\vec{e}_\mu, \tilde{e}^\nu$ chosen so that:

$$d\vec{x} = dx^\mu \vec{e}_\mu \text{ and}$$

$$df = \langle \tilde{d}f, d\vec{x} \rangle \text{ for any scalar field } f \text{ coordinate-free}$$

where $\tilde{d} = \tilde{e}^\mu \partial_\mu$ in a coordinate basis

$$\text{check: } df = \langle \tilde{d}f, d\vec{x} \rangle$$

$$= \langle \tilde{e}^\mu \partial_\mu f, dx^\nu \vec{e}_\nu \rangle$$

$$= \partial_\mu f dx^\nu \langle \tilde{e}^\mu, \vec{e}_\nu \rangle \text{ since scalars commute}$$

$$\text{i.e. } df = \partial_\mu f dx^\mu$$

GR: what is a coordinate basis?

coordinate basis: $\vec{e}_\mu, \tilde{e}^\nu$ chosen so that:

$$d\vec{x} = dx^\mu \vec{e}_\mu \text{ and}$$

$$df = \langle \tilde{d}f, d\vec{x} \rangle \text{ for any scalar field } f \text{ coordinate-free}$$

where $\tilde{d} = \tilde{e}^\mu \partial_\mu$ in a coordinate basis

$$\text{check: } df = \langle \tilde{d}f, d\vec{x} \rangle$$

$$= \langle \tilde{e}^\mu \partial_\mu f, dx^\nu \vec{e}_\nu \rangle$$

$$= \partial_\mu f dx^\nu \langle \tilde{e}^\mu, \vec{e}_\nu \rangle \text{ since scalars commute}$$

$$\text{i.e. } df = \partial_\mu f dx^\mu$$

$$\text{check: } \tilde{d}x^\mu = \tilde{e}^\nu \partial_\nu x^\mu$$

GR: what is a coordinate basis?

coordinate basis: $\vec{e}_\mu, \tilde{e}^\nu$ chosen so that:

$$d\vec{x} = dx^\mu \vec{e}_\mu \text{ and}$$

$$df = \langle \tilde{d}f, d\vec{x} \rangle \text{ for any scalar field } f \text{ coordinate-free}$$

where $\tilde{d} = \tilde{e}^\mu \partial_\mu$ in a coordinate basis

$$\text{check: } df = \langle \tilde{d}f, d\vec{x} \rangle$$

$$= \langle \tilde{e}^\mu \partial_\mu f, dx^\nu \vec{e}_\nu \rangle$$

$$= \partial_\mu f dx^\nu \langle \tilde{e}^\mu, \vec{e}_\nu \rangle \text{ since scalars commute}$$

$$\text{i.e. } df = \partial_\mu f dx^\mu$$

$$\text{check: } \tilde{d}x^\mu = \tilde{e}^\nu \partial_\nu x^\mu$$

$$= \sum_\nu \tilde{e}^\nu \partial_\nu x^\mu$$

GR: what is a coordinate basis?

coordinate basis: $\vec{e}_\mu, \tilde{e}^\nu$ chosen so that:

$$d\vec{x} = dx^\mu \vec{e}_\mu \text{ and}$$

$$df = \langle \tilde{d}f, d\vec{x} \rangle \text{ for any scalar field } f \text{ coordinate-free}$$

where $\tilde{d} = \tilde{e}^\mu \partial_\mu$ in a coordinate basis

$$\text{check: } df = \langle \tilde{d}f, d\vec{x} \rangle$$

$$= \langle \tilde{e}^\mu \partial_\mu f, dx^\nu \vec{e}_\nu \rangle$$

$$= \partial_\mu f dx^\nu \langle \tilde{e}^\mu, \vec{e}_\nu \rangle \text{ since scalars commute}$$

$$\text{i.e. } df = \partial_\mu f dx^\mu$$

$$\text{check: } \tilde{d}x^\mu = \tilde{e}^\nu \partial_\nu x^\mu$$

$$= \sum_\nu \tilde{e}^\nu \frac{\partial}{\partial x^\nu} x^\mu$$

GR: what is a coordinate basis?

coordinate basis: $\vec{e}_\mu, \tilde{e}^\nu$ chosen so that:

$$d\vec{x} = dx^\mu \vec{e}_\mu \text{ and}$$

$$df = \langle \tilde{d}f, d\vec{x} \rangle \text{ for any scalar field } f \text{ coordinate-free}$$

where $\tilde{d} = \tilde{e}^\mu \partial_\mu$ in a coordinate basis

$$\text{check: } df = \langle \tilde{d}f, d\vec{x} \rangle$$

$$= \langle \tilde{e}^\mu \partial_\mu f, dx^\nu \vec{e}_\nu \rangle$$

$$= \partial_\mu f dx^\nu \langle \tilde{e}^\mu, \vec{e}_\nu \rangle \text{ since scalars commute}$$

$$\text{i.e. } df = \partial_\mu f dx^\mu$$

$$\text{check: } \tilde{d}x^\mu = \tilde{e}^\nu \partial_\nu x^\mu$$

$$= \tilde{e}^\mu$$

GR: metric using $d\vec{x}$

we now have

GR: metric using $d\vec{x}$

we now have

$$ds^2 := |d\vec{x}|^2$$

GR: metric using $d\vec{x}$

we now have

$$ds^2 := |d\vec{x}|^2 = \mathbf{g}(d\vec{x}, d\vec{x})$$

GR: metric using $d\vec{x}$

we now have

$$ds^2 := |d\vec{x}|^2 = \mathbf{g}(d\vec{x}, d\vec{x}) = d\vec{x} \cdot d\vec{x} \text{ coordinate-free}$$

GR: metric using $d\vec{x}$

we now have

$$ds^2 := |d\vec{x}|^2 = \mathbf{g}(d\vec{x}, d\vec{x}) = d\vec{x} \cdot d\vec{x} \text{ coordinate-free}$$

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu \text{ if } x^\mu \text{ are a coordinate basis}$$

GR: e.g. Euclidean g on $\mathbb{R}^2$

$g_{r\theta}$ and g_{xy}

$$ds^2 = dx^2 + dy^2 = dr^2 + r^2 d\theta^2$$

GR: e.g. Euclidean g on $\mathbb{R}^2$

$g_{r\theta}$ and g_{xy}

$$ds^2 = dx^2 + dy^2 = dr^2 + r^2 d\theta^2$$

$$\vec{e}_x \cdot \vec{e}_x = 1 = \vec{e}_y \cdot \vec{e}_y, \text{ others zero}$$


GR: e.g. Euclidean g on $\mathbb{R}^2$

$g_{r\theta}$ and g_{xy}

$$ds^2 = dx^2 + dy^2 = dr^2 + r^2 d\theta^2$$

$$\vec{e}_x \cdot \vec{e}_x = 1 = \vec{e}_y \cdot \vec{e}_y, \text{ others zero}$$

$$\vec{e}_r \cdot \vec{e}_r = 1, \vec{e}_\theta \cdot \vec{e}_\theta = r^2 \neq 1$$



GR: e.g. Euclidean g on $\mathbb{R}^2$

$g_{r\theta}$ and g_{xy}

$$ds^2 = dx^2 + dy^2 = dr^2 + r^2 d\theta^2$$

$$\vec{e}_x \cdot \vec{e}_x = 1 = \vec{e}_y \cdot \vec{e}_y, \text{ others zero}$$

$$\vec{e}_r \cdot \vec{e}_r = 1, \vec{e}_\theta \cdot \vec{e}_\theta = r^2 \neq 1$$

$$g^{\mu\alpha} g_{\alpha\nu} = \delta^\mu_\nu \Rightarrow g^{xx} = 1 = g^{yy}, g^{xy} = 0 = g^{yx}$$

GR: e.g. Euclidean g on $\mathbb{R}^2$

$g_{r\theta}$ and g_{xy}

$$ds^2 = dx^2 + dy^2 = dr^2 + r^2 d\theta^2$$

$$\vec{e}_x \cdot \vec{e}_x = 1 = \vec{e}_y \cdot \vec{e}_y, \text{ others zero}$$

$$\vec{e}_r \cdot \vec{e}_r = 1, \vec{e}_\theta \cdot \vec{e}_\theta = r^2 \neq 1$$

$$g^{\mu\alpha} g_{\alpha\nu} = \delta^\mu_\nu \Rightarrow g^{xx} = 1 = g^{yy}, g^{xy} = 0 = g^{yx}$$

$$\text{but } g^{rr} = 1 \neq g^{\theta\theta} = r^{-2}, g^{r\theta} = 0 = g^{\theta r}$$

GR: e.g. Euclidean g on $\mathbb{R}^2$

$g_{r\theta}$ and g_{xy}


$$ds^2 = dx^2 + dy^2 = dr^2 + r^2 d\theta^2$$

$$\vec{e}_x \cdot \vec{e}_x = 1 = \vec{e}_y \cdot \vec{e}_y, \text{ others zero}$$

$$\vec{e}_r \cdot \vec{e}_r = 1, \vec{e}_\theta \cdot \vec{e}_\theta = r^2 \neq 1$$

$$g^{\mu\alpha} g_{\alpha\nu} = \delta^\mu_\nu \Rightarrow g^{xx} = 1 = g^{yy}, g^{xy} = 0 = g^{yx}$$

$$\text{but } g^{rr} = 1 \neq g^{\theta\theta} = r^{-2}, g^{r\theta} = 0 = g^{\theta r}$$



GR: e.g. Euclidean g on $\mathbb{R}^2$

$g_{r\theta}$ and g_{xy}


$$ds^2 = dx^2 + dy^2 = dr^2 + r^2 d\theta^2$$

$$\vec{e}_x \cdot \vec{e}_x = 1 = \vec{e}_y \cdot \vec{e}_y, \text{ others zero}$$

$$\vec{e}_r \cdot \vec{e}_r = 1, \vec{e}_\theta \cdot \vec{e}_\theta = r^2 \neq 1$$

$$g^{\mu\alpha} g_{\alpha\nu} = \delta^\mu_\nu \Rightarrow g^{xx} = 1 = g^{yy}, g^{xy} = 0 = g^{yx}$$

$$\text{but } g^{rr} = 1 \neq g^{\theta\theta} = r^{-2}, g^{r\theta} = 0 = g^{\theta r}$$



$$\text{so } \tilde{e}^r \cdot \tilde{e}^r = 1, \tilde{e}^\theta \cdot \tilde{e}^\theta = r^{-2} \neq 1$$

GR: gradient of a vector: $\nabla \vec{A}$

gradient of scalar field: $\tilde{d}\phi \equiv \tilde{\nabla}\phi$

GR: gradient of a vector: $\nabla \vec{A}$

what is gradient of vector field $\tilde{\nabla} \vec{A}$?

GR: gradient of a vector: $\nabla \vec{A}$

$$\tilde{\nabla} \vec{A} = \tilde{\nabla}(A^\nu \vec{e}_\nu)$$

GR: gradient of a vector: $\nabla \vec{A}$

$$\begin{aligned}\tilde{\nabla} \vec{A} &= \tilde{\nabla}(A^\nu \vec{e}_\nu) \\ &= \tilde{e}^\mu \partial_\mu (A^\nu \vec{e}_\nu)\end{aligned}$$

GR: gradient of a vector: $\nabla \vec{A}$

$$\begin{aligned}\tilde{\nabla} \vec{A} &= \tilde{\nabla}(A^\nu \vec{e}_\nu) \\ &= \tilde{e}^\mu \partial_\mu (A^\nu \vec{e}_\nu) \\ &= \tilde{e}^\mu \otimes [(\partial_\mu A^\nu) \vec{e}_\nu + A^\nu \partial_\mu \vec{e}_\nu] \text{ by product rule and linearity}\end{aligned}$$

GR: gradient of a vector: $\nabla \vec{A}$

$$\tilde{\nabla} \vec{A} = \tilde{\nabla}(A^\nu \vec{e}_\nu)$$

$$= \tilde{e}^\mu \partial_\mu (A^\nu \vec{e}_\nu)$$

$$= \partial_\mu A^\nu \tilde{e}^\mu \otimes \vec{e}_\nu + A^\nu \tilde{e}^\mu \otimes \partial_\mu \vec{e}_\nu$$

GR: gradient of a vector: $\nabla \vec{A}$

$$\tilde{\nabla} \vec{A} = \tilde{\nabla}(A^\nu \vec{e}_\nu)$$

$$= \tilde{e}^\mu \partial_\mu (A^\nu \vec{e}_\nu)$$

$$= \partial_\mu A^\nu \tilde{e}^\mu \otimes \vec{e}_\nu + A^\nu \tilde{e}^\mu \otimes \partial_\mu \vec{e}_\nu$$

give a name to the second part: it must be a linear combination of basis vectors $\vec{e}_\lambda$

GR: gradient of a vector: $\nabla \vec{A}$

$$\tilde{\nabla} \vec{A} = \tilde{\nabla}(A^\nu \vec{e}_\nu)$$

$$= \tilde{e}^\mu \partial_\mu (A^\nu \vec{e}_\nu)$$

$$= \partial_\mu A^\nu \tilde{e}^\mu \otimes \vec{e}_\nu + A^\nu \tilde{e}^\mu \otimes \partial_\mu \vec{e}_\nu$$

define $\Gamma_{\nu\mu}^\lambda \vec{e}_\lambda := \partial_\mu \vec{e}_\nu$ Christoffel symbols of second kind
(symmetric defn)

GR: gradient of a vector: $\nabla \vec{A}$

$$\tilde{\nabla} \vec{A} = \tilde{\nabla}(A^\nu \vec{e}_\nu)$$

$$= \tilde{e}^\mu \partial_\mu (A^\nu \vec{e}_\nu)$$

$$= \partial_\mu A^\nu \tilde{e}^\mu \otimes \vec{e}_\nu + A^\nu \tilde{e}^\mu \otimes \partial_\mu \vec{e}_\nu$$

define $\Gamma_{\nu\mu}^\lambda \vec{e}_\lambda := \partial_\mu \vec{e}_\nu$ Christoffel symbols of second kind
(symmetric defn)

$$\text{so } \tilde{\nabla} \vec{A} = \partial_\mu A^\nu \tilde{e}^\mu \otimes \vec{e}_\nu + A^\nu \tilde{e}^\mu \otimes \Gamma_{\nu\mu}^\lambda \vec{e}_\lambda$$

GR: gradient of a vector: $\nabla \vec{A}$

$$\tilde{\nabla} \vec{A} = \tilde{\nabla}(A^\nu \vec{e}_\nu)$$

$$= \tilde{e}^\mu \partial_\mu (A^\nu \vec{e}_\nu)$$

$$= \partial_\mu A^\nu \tilde{e}^\mu \otimes \vec{e}_\nu + A^\nu \tilde{e}^\mu \otimes \partial_\mu \vec{e}_\nu$$

define $\Gamma_{\nu\mu}^\lambda \vec{e}_\lambda := \partial_\mu \vec{e}_\nu$ Christoffel symbols of second kind
(symmetric defn)

$$\text{so } \tilde{\nabla} \vec{A} = \partial_\mu A^\nu \tilde{e}^\mu \otimes \vec{e}_\nu + A^\nu \tilde{e}^\mu \otimes \Gamma_{\nu\mu}^\lambda \vec{e}_\lambda$$

$$= \partial_\mu A^\nu \tilde{e}^\mu \otimes \vec{e}_\nu + A^\nu \Gamma_{\nu\mu}^\lambda \tilde{e}^\mu \otimes \vec{e}_\lambda \text{ since any } \Gamma_{\nu\mu}^\lambda \text{ is a scalar}$$

GR: gradient of a vector: $\nabla \vec{A}$

$$\tilde{\nabla} \vec{A} = \tilde{\nabla}(A^\nu \vec{e}_\nu)$$

$$= \tilde{e}^\mu \partial_\mu (A^\nu \vec{e}_\nu)$$

$$= \partial_\mu A^\nu \tilde{e}^\mu \otimes \vec{e}_\nu + A^\nu \tilde{e}^\mu \otimes \partial_\mu \vec{e}_\nu$$

define $\Gamma_{\nu\mu}^\lambda \vec{e}_\lambda := \partial_\mu \vec{e}_\nu$ Christoffel symbols of second kind
(symmetric defn)

$$\text{so } \tilde{\nabla} \vec{A} = \partial_\mu A^\nu \tilde{e}^\mu \otimes \vec{e}_\nu + A^\nu \tilde{e}^\mu \otimes \Gamma_{\nu\mu}^\lambda \vec{e}_\lambda$$

$$= \partial_\mu A^\nu \tilde{e}^\mu \otimes \vec{e}_\nu + A^\lambda \Gamma_{\lambda\mu}^\nu \tilde{e}^\mu \otimes \vec{e}_\nu$$

since name of summation index is arbitrary, e.g.

$$\sum_\lambda x^{-2\lambda} = \sum_\mu x^{-2\mu} = \sum_\nu x^{-2\nu}$$

GR: gradient of a vector: $\nabla \vec{A}$

$$\tilde{\nabla} \vec{A} = \tilde{\nabla}(A^\nu \vec{e}_\nu)$$

$$= \tilde{e}^\mu \partial_\mu (A^\nu \vec{e}_\nu)$$

$$= \partial_\mu A^\nu \tilde{e}^\mu \otimes \vec{e}_\nu + A^\nu \tilde{e}^\mu \otimes \partial_\mu \vec{e}_\nu$$

define $\Gamma_{\nu\mu}^\lambda \vec{e}_\lambda := \partial_\mu \vec{e}_\nu$ Christoffel symbols of second kind
(symmetric defn)

$$\text{so } \tilde{\nabla} \vec{A} = \partial_\mu A^\nu \tilde{e}^\mu \otimes \vec{e}_\nu + A^\nu \tilde{e}^\mu \otimes \Gamma_{\nu\mu}^\lambda \vec{e}_\lambda$$

$$= \partial_\mu A^\nu \tilde{e}^\mu \otimes \vec{e}_\nu + A^\lambda \Gamma_{\lambda\mu}^\nu \tilde{e}^\mu \otimes \vec{e}_\nu$$

$$= (\partial_\mu A^\nu + A^\lambda \Gamma_{\lambda\mu}^\nu) \tilde{e}^\mu \otimes \vec{e}_\nu$$

GR: gradient of a vector: $\nabla \vec{A}$

$$\tilde{\nabla} \vec{A} = \tilde{\nabla}(A^\nu \vec{e}_\nu)$$

$$= \tilde{e}^\mu \partial_\mu (A^\nu \vec{e}_\nu)$$

$$= \partial_\mu A^\nu \tilde{e}^\mu \otimes \vec{e}_\nu + A^\nu \tilde{e}^\mu \otimes \partial_\mu \vec{e}_\nu$$

define $\Gamma_{\nu\mu}^\lambda \vec{e}_\lambda := \partial_\mu \vec{e}_\nu$ Christoffel symbols of second kind
(symmetric defn)

$$\text{so } \tilde{\nabla} \vec{A} = \partial_\mu A^\nu \tilde{e}^\mu \otimes \vec{e}_\nu + A^\nu \tilde{e}^\mu \otimes \Gamma_{\nu\mu}^\lambda \vec{e}_\lambda$$

$$= \partial_\mu A^\nu \tilde{e}^\mu \otimes \vec{e}_\nu + A^\lambda \Gamma_{\lambda\mu}^\nu \tilde{e}^\mu \otimes \vec{e}_\nu$$

$$= (\partial_\mu A^\nu + A^\lambda \Gamma_{\lambda\mu}^\nu) \tilde{e}^\mu \otimes \vec{e}_\nu$$

$$\nabla_\mu A^\nu := A^\nu_{;\mu} := \partial_\mu A^\nu + A^\lambda \Gamma_{\lambda\mu}^\nu$$

w:covariant derivative of vector

GR: gradient of a vector: $\nabla \vec{A}$

$$\tilde{\nabla} \vec{A} = \tilde{\nabla}(A^\nu \vec{e}_\nu)$$

$$= \tilde{e}^\mu \partial_\mu (A^\nu \vec{e}_\nu)$$

$$= \partial_\mu A^\nu \tilde{e}^\mu \otimes \vec{e}_\nu + A^\nu \tilde{e}^\mu \otimes \partial_\mu \vec{e}_\nu$$

define $\Gamma_{\nu\mu}^\lambda \vec{e}_\lambda := \partial_\mu \vec{e}_\nu$ Christoffel symbols of second kind
(symmetric defn)

$$\text{so } \tilde{\nabla} \vec{A} = \partial_\mu A^\nu \tilde{e}^\mu \otimes \vec{e}_\nu + A^\nu \tilde{e}^\mu \otimes \Gamma_{\nu\mu}^\lambda \vec{e}_\lambda$$

$$= \partial_\mu A^\nu \tilde{e}^\mu \otimes \vec{e}_\nu + A^\lambda \Gamma_{\lambda\mu}^\nu \tilde{e}^\mu \otimes \vec{e}_\nu$$

$$= (\partial_\mu A^\nu + A^\lambda \Gamma_{\lambda\mu}^\nu) \tilde{e}^\mu \otimes \vec{e}_\nu$$

$$\nabla_\mu A^\nu := A^\nu_{;\mu} := A^\nu_{,\mu} + A^\lambda \Gamma_{\lambda\mu}^\nu$$

w:covariant derivative of vector

GR: $\tilde{\nabla}$ and $\Gamma^\nu_{\lambda\mu}$

mathematically deeper: $\tilde{\nabla}$, usually written just as ∇ , is the w:Levi-Civita connection

GR: $\tilde{\nabla}$ and $\Gamma^\nu_{\lambda\mu}$

mathematically deeper: $\tilde{\nabla}$, usually written just as ∇ , is the Levi-Civita connection

warning: $\Gamma^\nu_{\lambda\mu}$ are NOT the components of a tensor

GR: $\tilde{\nabla}$ and $\Gamma^\nu_{\lambda\mu}$

mathematically deeper: $\tilde{\nabla}$, usually written just as ∇ , is the w:Levi-Civita connection

warning: $\Gamma^\nu_{\lambda\mu}$ are NOT the components of a tensor

- $\tilde{\nabla}$ applied to a (m, n) -tensor field on a manifold gives an $(m, n + 1)$ -tensor field

GR: $\tilde{\nabla}$ and $\Gamma^\nu_{\lambda\mu}$

mathematically deeper: $\tilde{\nabla}$, usually written just as ∇ , is the Levi-Civita connection

warning: $\Gamma^\nu_{\lambda\mu}$ are NOT the components of a tensor

- $\tilde{\nabla}$ applied to a (m, n) -tensor field on a manifold gives an $(m, n + 1)$ -tensor field

so far we showed how $\tilde{\nabla}$ applied to a $(0, 0)$ -tensor field = scalar field ϕ gives a $(0, 1)$ -tensor field = one-form field = $(\tilde{d}\phi)_\mu \vec{e}^\mu$

GR: $\tilde{\nabla}$ and $\Gamma^\nu_{\lambda\mu}$

mathematically deeper: $\tilde{\nabla}$, usually written just as ∇ , is the Levi-Civita connection

warning: $\Gamma^\nu_{\lambda\mu}$ are NOT the components of a tensor

- $\tilde{\nabla}$ applied to a (m, n) -tensor field on a manifold gives an $(m, n + 1)$ -tensor field

so far we showed how $\tilde{\nabla}$ applied to a $(0, 0)$ -tensor field = scalar field ϕ gives a $(0, 1)$ -tensor field = one-form field =

$$\tilde{\nabla}_\mu \phi \vec{e}^\mu = \partial_\mu \phi \vec{e}^\mu$$

GR: $\tilde{\nabla}$ and $\Gamma^\nu_{\lambda\mu}$

mathematically deeper: $\tilde{\nabla}$, usually written just as ∇ , is the w:Levi-Civita connection

warning: $\Gamma^\nu_{\lambda\mu}$ are NOT the components of a tensor

- $\tilde{\nabla}$ applied to a (m, n) -tensor field on a manifold gives an $(m, n + 1)$ -tensor field

so far we showed how $\tilde{\nabla}$ applied to a $(0, 0)$ -tensor field = scalar field ϕ gives a $(0, 1)$ -tensor field = one-form field =

$$\tilde{\nabla}_\mu \phi \vec{e}^\mu = \partial_\mu \phi \vec{e}^\mu$$

and $\tilde{\nabla}$ on a $(1, 0)$ -tensor field = vector field $\vec{A}$ gives a $(1, 1)$ -tensor with components $\nabla_\mu A^\nu = \partial_\mu A^\nu + A^\lambda \Gamma^\nu_{\lambda\mu}$

GR: $\tilde{\nabla}$ and $\Gamma^\nu_{\lambda\mu}$

mathematically deeper: $\tilde{\nabla}$, usually written just as ∇ , is the w:Levi-Civita connection

warning: $\Gamma^\nu_{\lambda\mu}$ are NOT the components of a tensor

- $\tilde{\nabla}$ applied to a (m, n) -tensor field on a manifold gives an $(m, n + 1)$ -tensor field

so far we showed how $\tilde{\nabla}$ applied to a $(0, 0)$ -tensor field = scalar field ϕ gives a $(0, 1)$ -tensor field = one-form field =

$$\tilde{\nabla}_\mu \phi \vec{e}^\mu = \partial_\mu \phi \vec{e}^\mu$$

and $\tilde{\nabla}$ on a $(1, 0)$ -tensor field = vector field $\vec{A}$ gives a $(1, 1)$ -tensor with components $\nabla_\mu A^\nu = \partial_\mu A^\nu + A^\lambda \Gamma^\nu_{\lambda\mu}$

- tensors: $\tilde{\nabla} \phi = \tilde{\nabla}_\mu \phi \vec{e}^\mu$,

GR: $\tilde{\nabla}$ and $\Gamma^\nu_{\lambda\mu}$

mathematically deeper: $\tilde{\nabla}$, usually written just as ∇ , is the w:Levi-Civita connection

warning: $\Gamma^\nu_{\lambda\mu}$ are NOT the components of a tensor

- $\tilde{\nabla}$ applied to a (m, n) -tensor field on a manifold gives an $(m, n + 1)$ -tensor field

so far we showed how $\tilde{\nabla}$ applied to a $(0, 0)$ -tensor field = scalar field ϕ gives a $(0, 1)$ -tensor field = one-form field =

$$\tilde{\nabla}_\mu \phi \vec{e}^\mu = \partial_\mu \phi \vec{e}^\mu$$

and $\tilde{\nabla}$ on a $(1, 0)$ -tensor field = vector field $\vec{A}$ gives a $(1, 1)$ -tensor with components $\nabla_\mu A^\nu = \partial_\mu A^\nu + A^\lambda \Gamma^\nu_{\lambda\mu}$

- tensors: $\tilde{\nabla} \phi = \tilde{\nabla}_\mu \phi \vec{e}^\mu$, $\tilde{\nabla} \vec{A} = \left(\partial_\mu A^\nu + A^\lambda \Gamma^\nu_{\lambda\mu} \right) \vec{e}^\mu \otimes \vec{e}^\nu$

GR: $\tilde{\nabla}$ and $\Gamma^\nu_{\lambda\mu}$

mathematically deeper: $\tilde{\nabla}$, usually written just as ∇ , is the w:Levi-Civita connection

warning: $\Gamma^\nu_{\lambda\mu}$ are NOT the components of a tensor

- $\tilde{\nabla}$ applied to a (m, n) -tensor field on a manifold gives an $(m, n + 1)$ -tensor field

so far we showed how $\tilde{\nabla}$ applied to a $(0, 0)$ -tensor field = scalar field ϕ gives a $(0, 1)$ -tensor field = one-form field =

$$\tilde{\nabla}_\mu \phi \vec{e}^\mu = \partial_\mu \phi \vec{e}^\mu$$

and $\tilde{\nabla}$ on a $(1, 0)$ -tensor field = vector field $\vec{A}$ gives a $(1, 1)$ -tensor with components $\nabla_\mu A^\nu = \partial_\mu A^\nu + A^\lambda \Gamma^\nu_{\lambda\mu}$

- tensors: $\tilde{\nabla} \phi = \tilde{\nabla}_\mu \phi \vec{e}^\mu$, $\tilde{\nabla} \vec{A} = \left(\partial_\mu A^\nu + A^\lambda \Gamma^\nu_{\lambda\mu} \right) \vec{e}^\mu \otimes \vec{e}^\nu$

- not components of tensor: $\Gamma^\nu_{\lambda\mu}$

GR: gradient of one-form $\tilde{\nabla} \tilde{A}$

how does a one-form change with position? $\tilde{\nabla} \tilde{A} = ?$

GR: gradient of one-form $\tilde{\nabla} \tilde{A}$

evaluating $\tilde{\nabla} \tilde{A}$ as we did $\tilde{\nabla} \vec{A}$ shows that we again need $\partial_\mu \tilde{e}^\nu = F^\nu_{\lambda\mu} \tilde{e}^\lambda$ for some coefficients $F^\nu_{\lambda\mu}$

GR: gradient of one-form $\tilde{\nabla} \tilde{A}$

evaluating $\tilde{\nabla} \tilde{A}$ as we did $\tilde{\nabla} \vec{A}$ shows that we again need $\partial_\mu \tilde{e}^\nu = F^\nu_{\lambda\mu} \tilde{e}^\lambda$ for some coefficients $F^\nu_{\lambda\mu}$

how can we relate $\Gamma^\nu_{\lambda\mu}$ to $F^\nu_{\lambda\mu}$?

GR: gradient of one-form $\tilde{\nabla} \tilde{A}$

evaluating $\tilde{\nabla} \tilde{A}$ as we did $\tilde{\nabla} \vec{A}$ shows that we again need $\partial_\mu \tilde{e}^\nu = F^\nu_{\lambda\mu} \tilde{e}^\lambda$ for some coefficients $F^\nu_{\lambda\mu}$

how can we relate $\Gamma^\nu_{\lambda\mu}$ to $F^\nu_{\lambda\mu}$?

relation between vectors and one-forms: $\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle = \delta^\nu_\lambda$

GR: gradient of one-form $\tilde{\nabla} \tilde{A}$

evaluating $\tilde{\nabla} \tilde{A}$ as we did $\tilde{\nabla} \vec{A}$ shows that we again need $\partial_\mu \tilde{e}^\nu = F^\nu_{\lambda\mu} \tilde{e}^\lambda$ for some coefficients $F^\nu_{\lambda\mu}$

how can we relate $\Gamma^\nu_{\lambda\mu}$ to $F^\nu_{\lambda\mu}$?

relation between vectors and one-forms: $\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle = \delta^\nu_\lambda$

$\partial_\mu \delta^\nu_\lambda = 0$ (obviously)

GR: gradient of one-form $\tilde{\nabla} \tilde{A}$

evaluating $\tilde{\nabla} \tilde{A}$ as we did $\tilde{\nabla} \vec{A}$ shows that we again need $\partial_\mu \tilde{e}^\nu = F^\nu_{\lambda\mu} \tilde{e}^\lambda$ for some coefficients $F^\nu_{\lambda\mu}$

how can we relate $\Gamma^\nu_{\lambda\mu}$ to $F^\nu_{\lambda\mu}$?

relation between vectors and one-forms: $\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle = \delta^\nu_\lambda$

$$0 = \partial_\mu \delta^\nu_\lambda = \partial_\mu (\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle)$$

GR: gradient of one-form $\tilde{\nabla} \tilde{A}$

evaluating $\tilde{\nabla} \tilde{A}$ as we did $\tilde{\nabla} \vec{A}$ shows that we again need $\partial_\mu \tilde{e}^\nu = F^\nu_{\lambda\mu} \tilde{e}^\lambda$ for some coefficients $F^\nu_{\lambda\mu}$

how can we relate $\Gamma^\nu_{\lambda\mu}$ to $F^\nu_{\lambda\mu}$?

relation between vectors and one-forms: $\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle = \delta^\nu_\lambda$

$$0 = \partial_\mu \delta^\nu_\lambda = \partial_\mu (\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle)$$

can we use the product rule with this scalar product?

$$\partial_\mu (\langle \tilde{A}, \vec{B} \rangle) = ?$$

GR: gradient of one-form $\tilde{\nabla} \tilde{A}$

evaluating $\tilde{\nabla} \tilde{A}$ as we did $\tilde{\nabla} \vec{A}$ shows that we again need $\partial_\mu \tilde{e}^\nu = F^\nu_{\lambda\mu} \tilde{e}^\lambda$ for some coefficients $F^\nu_{\lambda\mu}$

how can we relate $\Gamma^\nu_{\lambda\mu}$ to $F^\nu_{\lambda\mu}$?

relation between vectors and one-forms: $\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle = \delta^\nu_\lambda$

$$0 = \partial_\mu \delta^\nu_\lambda = \partial_\mu (\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle)$$

can we use the product rule with this scalar product?

$$\partial_\mu \left(\langle \tilde{A}, \vec{B} \rangle \right) = \partial_\mu (A_\nu B^\nu) \text{ in some coordinate basis}$$

GR: gradient of one-form $\tilde{\nabla} \tilde{A}$

evaluating $\tilde{\nabla} \tilde{A}$ as we did $\tilde{\nabla} \vec{A}$ shows that we again need $\partial_\mu \tilde{e}^\nu = F^\nu_{\lambda\mu} \tilde{e}^\lambda$ for some coefficients $F^\nu_{\lambda\mu}$

how can we relate $\Gamma^\nu_{\lambda\mu}$ to $F^\nu_{\lambda\mu}$?

relation between vectors and one-forms: $\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle = \delta^\nu_\lambda$

$$0 = \partial_\mu \delta^\nu_\lambda = \partial_\mu (\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle)$$

can we use the product rule with this scalar product?

$$\partial_\mu \left(\langle \tilde{A}, \vec{B} \rangle \right) = \partial_\mu (A_\nu B^\nu)$$

$$= (\partial_\mu A_\nu) B^\nu + A_\nu (\partial_\mu B^\nu) \text{ by product rule on functions}$$

GR: gradient of one-form $\tilde{\nabla} \tilde{A}$

evaluating $\tilde{\nabla} \tilde{A}$ as we did $\tilde{\nabla} \vec{A}$ shows that we again need $\partial_\mu \tilde{e}^\nu = F^\nu_{\lambda\mu} \tilde{e}^\lambda$ for some coefficients $F^\nu_{\lambda\mu}$

how can we relate $\Gamma^\nu_{\lambda\mu}$ to $F^\nu_{\lambda\mu}$?

relation between vectors and one-forms: $\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle = \delta^\nu_\lambda$

$$0 = \partial_\mu \delta^\nu_\lambda = \partial_\mu (\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle)$$

can we use the product rule with this scalar product?

$$\partial_\mu \left(\langle \tilde{A}, \vec{B} \rangle \right) = \partial_\mu (A_\nu B^\nu)$$

$$= (\partial_\mu A_\nu) B^\nu + A_\nu (\partial_\mu B^\nu)$$

$$= \langle \partial_\mu \tilde{A}, \vec{B} \rangle + \langle \tilde{A}, \partial_\mu \vec{B} \rangle \text{ since}$$

$$\partial_\mu \tilde{A} = (\partial_\mu A_0, \partial_\mu A_1, \partial_\mu A_2, \partial_\mu A_3)$$

GR: gradient of one-form $\tilde{\nabla} \tilde{A}$

evaluating $\tilde{\nabla} \tilde{A}$ as we did $\tilde{\nabla} \vec{A}$ shows that we again need $\partial_\mu \tilde{e}^\nu = F^\nu_{\lambda\mu} \tilde{e}^\lambda$ for some coefficients $F^\nu_{\lambda\mu}$

how can we relate $\Gamma^\nu_{\lambda\mu}$ to $F^\nu_{\lambda\mu}$?

relation between vectors and one-forms: $\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle = \delta^\nu_\lambda$

$$0 = \partial_\mu \delta^\nu_\lambda = \partial_\mu (\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle)$$

$$\text{product rule holds: } \partial_\mu \left(\langle \tilde{A}, \vec{B} \rangle \right) = \langle \partial_\mu \tilde{A}, \vec{B} \rangle + \langle \tilde{A}, \partial_\mu \vec{B} \rangle$$

GR: gradient of one-form $\tilde{\nabla} \tilde{A}$

evaluating $\tilde{\nabla} \tilde{A}$ as we did $\tilde{\nabla} \vec{A}$ shows that we again need $\partial_\mu \tilde{e}^\nu = F^\nu_{\lambda\mu} \tilde{e}^\lambda$ for some coefficients $F^\nu_{\lambda\mu}$

how can we relate $\Gamma^\nu_{\lambda\mu}$ to $F^\nu_{\lambda\mu}$?

relation between vectors and one-forms: $\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle = \delta^\nu_\lambda$

$$0 = \partial_\mu \delta^\nu_\lambda = \partial_\mu (\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle)$$

$$\text{product rule holds: } \partial_\mu (\langle \tilde{A}, \vec{B} \rangle) = \langle \partial_\mu \tilde{A}, \vec{B} \rangle + \langle \tilde{A}, \partial_\mu \vec{B} \rangle$$

$$\text{so } 0 = \langle \partial_\mu \tilde{e}^\nu, \vec{e}_\lambda \rangle + \langle \tilde{e}^\nu, \partial_\mu \vec{e}_\lambda \rangle$$

GR: gradient of one-form $\tilde{\nabla} \tilde{A}$

evaluating $\tilde{\nabla} \tilde{A}$ as we did $\tilde{\nabla} \vec{A}$ shows that we again need $\partial_\mu \tilde{e}^\nu = F^\nu_{\lambda\mu} \tilde{e}^\lambda$ for some coefficients $F^\nu_{\lambda\mu}$

how can we relate $\Gamma^\nu_{\lambda\mu}$ to $F^\nu_{\lambda\mu}$?

relation between vectors and one-forms: $\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle = \delta^\nu_\lambda$

$$0 = \partial_\mu \delta^\nu_\lambda = \partial_\mu (\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle)$$

$$\text{product rule holds: } \partial_\mu (\langle \tilde{A}, \vec{B} \rangle) = \langle \partial_\mu \tilde{A}, \vec{B} \rangle + \langle \tilde{A}, \partial_\mu \vec{B} \rangle$$

$$\text{so } 0 = \langle \partial_\mu \tilde{e}^\nu, \vec{e}_\lambda \rangle + \langle \tilde{e}^\nu, \partial_\mu \vec{e}_\lambda \rangle$$

$$= \langle F^\nu_{\kappa\mu} \tilde{e}^\kappa, \vec{e}_\lambda \rangle + \langle \tilde{e}^\nu, \Gamma^\kappa_{\lambda\mu} \vec{e}_\kappa \rangle$$

GR: gradient of one-form $\tilde{\nabla} \tilde{A}$

evaluating $\tilde{\nabla} \tilde{A}$ as we did $\tilde{\nabla} \vec{A}$ shows that we again need $\partial_\mu \tilde{e}^\nu = F^\nu_{\lambda\mu} \tilde{e}^\lambda$ for some coefficients $F^\nu_{\lambda\mu}$

how can we relate $\Gamma^\nu_{\lambda\mu}$ to $F^\nu_{\lambda\mu}$?

relation between vectors and one-forms: $\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle = \delta^\nu_\lambda$

$$0 = \partial_\mu \delta^\nu_\lambda = \partial_\mu (\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle)$$

$$\text{product rule holds: } \partial_\mu \left(\langle \tilde{A}, \vec{B} \rangle \right) = \langle \partial_\mu \tilde{A}, \vec{B} \rangle + \langle \tilde{A}, \partial_\mu \vec{B} \rangle$$

$$\text{so } 0 = \langle \partial_\mu \tilde{e}^\nu, \vec{e}_\lambda \rangle + \langle \tilde{e}^\nu, \partial_\mu \vec{e}_\lambda \rangle$$

$$= F^\nu_{\kappa\mu} \langle \tilde{e}^\kappa, \vec{e}_\lambda \rangle + \Gamma^\kappa_{\lambda\mu} \langle \tilde{e}^\nu, \vec{e}_\kappa \rangle$$

GR: gradient of one-form $\tilde{\nabla} \tilde{A}$

evaluating $\tilde{\nabla} \tilde{A}$ as we did $\tilde{\nabla} \vec{A}$ shows that we again need $\partial_\mu \tilde{e}^\nu = F^\nu_{\lambda\mu} \tilde{e}^\lambda$ for some coefficients $F^\nu_{\lambda\mu}$

how can we relate $\Gamma^\nu_{\lambda\mu}$ to $F^\nu_{\lambda\mu}$?

relation between vectors and one-forms: $\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle = \delta^\nu_\lambda$

$$0 = \partial_\mu \delta^\nu_\lambda = \partial_\mu (\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle)$$

$$\text{product rule holds: } \partial_\mu (\langle \tilde{A}, \vec{B} \rangle) = \langle \partial_\mu \tilde{A}, \vec{B} \rangle + \langle \tilde{A}, \partial_\mu \vec{B} \rangle$$

$$\begin{aligned} \text{so } 0 &= \langle \partial_\mu \tilde{e}^\nu, \vec{e}_\lambda \rangle + \langle \tilde{e}^\nu, \partial_\mu \vec{e}_\lambda \rangle \\ &= F^\nu_{\lambda\mu} + \Gamma^\nu_{\lambda\mu} \text{ since } \langle \tilde{e}^\kappa, \vec{e}_\lambda \rangle = \delta^\kappa_\lambda \end{aligned}$$

GR: gradient of one-form $\tilde{\nabla} \tilde{A}$

evaluating $\tilde{\nabla} \tilde{A}$ as we did $\tilde{\nabla} \vec{A}$ shows that we again need $\partial_\mu \tilde{e}^\nu = F^\nu_{\lambda\mu} \tilde{e}^\lambda$ for some coefficients $F^\nu_{\lambda\mu}$

how can we relate $\Gamma^\nu_{\lambda\mu}$ to $F^\nu_{\lambda\mu}$?

relation between vectors and one-forms: $\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle = \delta^\nu_\lambda$

$$0 = \partial_\mu \delta^\nu_\lambda = \partial_\mu (\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle)$$

$$\text{product rule holds: } \partial_\mu (\langle \tilde{A}, \vec{B} \rangle) = \langle \partial_\mu \tilde{A}, \vec{B} \rangle + \langle \tilde{A}, \partial_\mu \vec{B} \rangle$$

$$\text{so } 0 = \langle \partial_\mu \tilde{e}^\nu, \vec{e}_\lambda \rangle + \langle \tilde{e}^\nu, \partial_\mu \vec{e}_\lambda \rangle$$

$$= F^\nu_{\lambda\mu} + \Gamma^\nu_{\lambda\mu} \text{ since } \langle \tilde{e}^\kappa, \vec{e}_\lambda \rangle = \delta^\kappa_\lambda$$

$$\text{hence, } \partial_\mu \tilde{e}^\nu =: F^\nu_{\lambda\mu} \tilde{e}^\lambda = -\Gamma^\nu_{\lambda\mu} \tilde{e}^\lambda$$

GR: gradient of one-form $\tilde{\nabla} \tilde{A}$

evaluating $\tilde{\nabla} \tilde{A}$ as we did $\tilde{\nabla} \vec{A}$ shows that we again need $\partial_\mu \tilde{e}^\nu = F^\nu_{\lambda\mu} \tilde{e}^\lambda$ for some coefficients $F^\nu_{\lambda\mu}$

how can we relate $\Gamma^\nu_{\lambda\mu}$ to $F^\nu_{\lambda\mu}$?

relation between vectors and one-forms: $\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle = \delta^\nu_\lambda$

$$0 = \partial_\mu \delta^\nu_\lambda = \partial_\mu (\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle)$$

$$\text{product rule holds: } \partial_\mu (\langle \tilde{A}, \vec{B} \rangle) = \langle \partial_\mu \tilde{A}, \vec{B} \rangle + \langle \tilde{A}, \partial_\mu \vec{B} \rangle$$

$$\text{so } 0 = \langle \partial_\mu \tilde{e}^\nu, \vec{e}_\lambda \rangle + \langle \tilde{e}^\nu, \partial_\mu \vec{e}_\lambda \rangle$$

$$= F^\nu_{\lambda\mu} + \Gamma^\nu_{\lambda\mu} \text{ since } \langle \tilde{e}^\kappa, \vec{e}_\lambda \rangle = \delta^\kappa_\lambda$$

$$\text{hence, } \partial_\mu \tilde{e}^\nu =: F^\nu_{\lambda\mu} \tilde{e}^\lambda = -\Gamma^\nu_{\lambda\mu} \tilde{e}^\lambda$$

$$\tilde{\nabla}_\mu A^\nu = \partial_\mu A^\nu + A^\lambda \Gamma^\nu_{\lambda\mu} \quad , \quad \tilde{\nabla}_\mu A_\nu = \partial_\mu A_\nu - A_\lambda \Gamma^\lambda_{\mu\nu}$$

GR: gradient of one-form $\tilde{\nabla} \tilde{A}$

evaluating $\tilde{\nabla} \tilde{A}$ as we did $\tilde{\nabla} \vec{A}$ shows that we again need $\partial_\mu \tilde{e}^\nu = F^\nu_{\lambda\mu} \tilde{e}^\lambda$ for some coefficients $F^\nu_{\lambda\mu}$

how can we relate $\Gamma^\nu_{\lambda\mu}$ to $F^\nu_{\lambda\mu}$?

relation between vectors and one-forms: $\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle = \delta^\nu_\lambda$

$$0 = \partial_\mu \delta^\nu_\lambda = \partial_\mu (\langle \tilde{e}^\nu, \vec{e}_\lambda \rangle)$$

$$\text{product rule holds: } \partial_\mu (\langle \tilde{A}, \vec{B} \rangle) = \langle \partial_\mu \tilde{A}, \vec{B} \rangle + \langle \tilde{A}, \partial_\mu \vec{B} \rangle$$

$$\text{so } 0 = \langle \partial_\mu \tilde{e}^\nu, \vec{e}_\lambda \rangle + \langle \tilde{e}^\nu, \partial_\mu \vec{e}_\lambda \rangle$$

$$= F^\nu_{\lambda\mu} + \Gamma^\nu_{\lambda\mu} \text{ since } \langle \tilde{e}^\kappa, \vec{e}_\lambda \rangle = \delta^\kappa_\lambda$$

$$\text{hence, } \partial_\mu \tilde{e}^\nu =: F^\nu_{\lambda\mu} \tilde{e}^\lambda = -\Gamma^\nu_{\lambda\mu} \tilde{e}^\lambda$$

$$A^\nu_{;\mu} = A^\nu_{,\mu} + A^\lambda \Gamma^\nu_{\lambda\mu} \quad , \quad A_{\nu;\mu} = A_{\nu,\mu} - A_\lambda \Gamma^\lambda_{\mu\nu}$$

GR: smooth manifold and $\tilde{\nabla}g$

similarly, we can write the $(0, 3)$ -tensor

$$\tilde{\nabla}g = (\nabla_{\lambda}g_{\mu\nu})\tilde{e}^{\lambda} \otimes \tilde{e}^{\mu} \otimes \tilde{e}^{\nu}$$

$$\text{giving } \nabla_{\lambda}g_{\mu\nu} = \partial_{\lambda}g_{\mu\nu} - \Gamma^{\kappa}_{\mu\lambda}g_{\kappa\nu} - \Gamma^{\kappa}_{\nu\lambda}g_{\mu\kappa}$$

GR: smooth manifold and $\tilde{\nabla}g$

similarly, we can write the $(0, 3)$ -tensor

$$\tilde{\nabla}g = (\nabla_\lambda g_{\mu\nu}) \tilde{e}^\lambda \otimes \tilde{e}^\mu \otimes \tilde{e}^\nu$$

$$\text{giving } \nabla_\lambda g_{\mu\nu} = \partial_\lambda g_{\mu\nu} - \Gamma_{\mu\lambda}^\kappa g_{\kappa\nu} - \Gamma_{\nu\lambda}^\kappa g_{\mu\kappa}$$

$$\text{also } \tilde{\nabla}g^{-1} = (\nabla_\lambda g^{\mu\nu}) \tilde{e}^\lambda \otimes \tilde{e}_\mu \otimes \tilde{e}_\nu$$

$$\text{and } \nabla_\lambda g^{\mu\nu} = \partial_\lambda g^{\mu\nu} + \Gamma_{\kappa\lambda}^\mu g^{\kappa\nu} + \Gamma_{\kappa\lambda}^\nu g^{\mu\kappa}$$

GR: smooth manifold and $\tilde{\nabla}g$

similarly, we can write the $(0, 3)$ -tensor

$$\tilde{\nabla}g = (\nabla_\lambda g_{\mu\nu}) \tilde{e}^\lambda \otimes \tilde{e}^\mu \otimes \tilde{e}^\nu$$

$$\text{giving } \nabla_\lambda g_{\mu\nu} = \partial_\lambda g_{\mu\nu} - \Gamma_{\mu\lambda}^\kappa g_{\kappa\nu} - \Gamma_{\nu\lambda}^\kappa g_{\mu\kappa}$$

$$\text{also } \tilde{\nabla}g^{-1} = (\nabla_\lambda g^{\mu\nu}) \tilde{e}^\lambda \otimes \tilde{e}_\mu \otimes \tilde{e}_\nu$$

$$\text{and } \nabla_\lambda g^{\mu\nu} = \partial_\lambda g^{\mu\nu} + \Gamma_{\kappa\lambda}^\mu g^{\kappa\nu} + \Gamma_{\kappa\lambda}^\nu g^{\mu\kappa}$$

Do we know anything interesting about $\tilde{\nabla}g$ for the manifolds of interest to GR?

GR: smooth manifold and $\tilde{\nabla}g$

similarly, we can write the $(0, 3)$ -tensor

$$\tilde{\nabla}g = (\nabla_{\lambda}g_{\mu\nu})\tilde{e}^{\lambda} \otimes \tilde{e}^{\mu} \otimes \tilde{e}^{\nu}$$

$$\text{giving } \nabla_{\lambda}g_{\mu\nu} = \partial_{\lambda}g_{\mu\nu} - \Gamma^{\kappa}_{\mu\lambda}g_{\kappa\nu} - \Gamma^{\kappa}_{\nu\lambda}g_{\mu\kappa}$$

$$\text{also } \tilde{\nabla}g^{-1} = (\nabla_{\lambda}g^{\mu\nu})\tilde{e}^{\lambda} \otimes \tilde{e}_{\mu} \otimes \tilde{e}_{\nu}$$

$$\text{and } \nabla_{\lambda}g^{\mu\nu} = \partial_{\lambda}g^{\mu\nu} + \Gamma^{\mu}_{\kappa\lambda}g^{\kappa\nu} + \Gamma^{\nu}_{\kappa\lambda}g^{\mu\kappa}$$

Do we know anything interesting about $\tilde{\nabla}g$ for the manifolds of interest to GR?

First, we need a rough description of the manifolds we need for GR.

GR: smooth manifold and $\tilde{\nabla}g$

topological 4-(pseudo-)manifold M
w:Manifold#Mathematical_definition

- only topological properties needed

GR: smooth manifold and $\tilde{\nabla}g$

topological 4-(pseudo-)manifold M
w:Manifold#Mathematical_definition

- only topological properties needed
- no differentiability, no metric needed

GR: smooth manifold and $\tilde{\nabla}g$


topological 4-(pseudo-)manifold M
w:Manifold#Mathematical_definition

- only topological properties needed
- next: relation with $\mathbb{R}^4$ (or M^4)

GR: smooth manifold and $\tilde{\nabla}g$

topological 4-(pseudo-)manifold M
w:Manifold#Mathematical_definition

- only topological properties needed
- next: relation with $\mathbb{R}^4$ (or M^4)



GR: smooth manifold and $\tilde{\nabla}g$


topological 4-(pseudo-)manifold M
w:Manifold#Mathematical_definition

- only topological properties needed

next: relation with $\mathbb{R}^4$ (or M^4)

w:Manifold

- chart := function ϕ_α from part of pseudo-4-manifold M to part of M^4 (Minkowski)
- atlas := set of overlapping charts that cover M




GR: smooth manifold and $\tilde{\nabla}g$

if every transition chart $:= \phi_\beta \circ \phi_\alpha^{-1}$ in an atlas for M is differentiable on $\mathbb{R}^4$ (or M^4), then M is a w:differentiable 4-(pseudo-)manifold

GR: smooth manifold and $\tilde{\nabla}g$


if every transition chart $:= \phi_\beta \circ \phi_\alpha^{-1}$ in an atlas for M is differentiable on $\mathbb{R}^4$ (or M^4), then M is a w:differentiable 4-(pseudo-)manifold



W:

GR: smooth manifold and $\tilde{\nabla}g$

if every transition chart $:= \phi_\beta \circ \phi_\alpha^{-1}$ in an atlas for M is differentiable on $\mathbb{R}^4$ (or M^4), then M is a w:differentiable 4-(pseudo-)manifold




w:

projections (left-to-right) ϕ_1, ϕ_2, ϕ_3 from S^2 to $\mathbb{R}^2$

GR: smooth manifold and $\tilde{\nabla}g$

if every transition chart $:= \phi_\beta \circ \phi_\alpha^{-1}$ in an atlas for M is differentiable on $\mathbb{R}^4$ (or M^4), then M is a w:differentiable 4-(pseudo-)manifold




w:

ϕ_1 is not differentiable, so $\phi_1 \circ \phi_2^{-1}$ is not differentiable

GR: smooth manifold and $\tilde{\nabla}g$

if every transition chart $:= \phi_\beta \circ \phi_\alpha^{-1}$ in an atlas for M is differentiable on $\mathbb{R}^4$ (or M^4), then M is a w:differentiable 4-(pseudo-)manifold



w:

atlas not enough to show that $S^2 =$ differentiable
2-manifold

GR: smooth manifold and $\tilde{\nabla}g$

if every transition chart $:= \phi_\beta \circ \phi_\alpha^{-1}$ in an atlas for M is differentiable on $\mathbb{R}^4$ (or M^4), then M is a w:differentiable 4-(pseudo-)manifold

GR: smooth manifold and $\tilde{\nabla}g$

if every transition chart $:= \phi_\beta \circ \phi_\alpha^{-1}$ in an atlas for M is differentiable on $\mathbb{R}^4$ (or M^4), then M is a w:differentiable 4-(pseudo-)manifold

if $\forall k \geq 1, \exists k$ -th derivatives, then M is a smooth 4-(pseudo-)manifold

GR: smooth manifold and $\tilde{\nabla}g$

if every transition chart $:= \phi_\beta \circ \phi_\alpha^{-1}$ in an atlas for M is differentiable on $\mathbb{R}^4$ (or M^4), then M is a w:differentiable 4-(pseudo-)manifold

if $\forall k \geq 1, \exists k$ -th derivatives, then M is a smooth 4-(pseudo-)manifold

if a (pseudo-)w:Riemannian metric g can be added to M , then (M, g) is a (pseudo-)Riemannian 4-manifold

GR: smooth manifold and $\tilde{\nabla}g$

if every transition chart $:= \phi_\beta \circ \phi_\alpha^{-1}$ in an atlas for M is differentiable on $\mathbb{R}^4$ (or M^4), then M is a w:differentiable 4-(pseudo-)manifold

if $\forall k \geq 1, \exists k$ -th derivatives, then M is a smooth 4-(pseudo-)manifold

if a (pseudo-)w:Riemannian metric g can be added to M , then (M, g) is a (pseudo-)Riemannian 4-manifold

if g has signature $(1, n-1)$ (i.e. $(-, +, +, +)$ or $(+, -, -, -)$, etc.), then (M, g) is a Lorentzian n -manifold

GR: smooth manifold and $\tilde{\nabla}g$

topological manifolds

GR: smooth manifold and $\tilde{\nabla}g$

topological manifolds

differentiable (pseudo-)manifolds

GR: smooth manifold and $\tilde{\nabla}g$



topological manifolds

differentiable (pseudo-)manifolds

smooth (pseudo-)manifolds



GR: smooth manifold and $\tilde{\nabla}g$



topological manifolds

differentiable (pseudo-)manifolds

smooth (pseudo-)manifolds

(pseudo-)Riemannian manifolds



GR: smooth manifold and $\tilde{\nabla}g$



topological manifolds

differentiable (pseudo-)manifolds

smooth (pseudo-)manifolds

(pseudo-)Riemannian manifolds

Lorentzian manifolds



GR: smooth manifold and $\tilde{\nabla}g$



topological manifolds

differentiable (pseudo-)manifolds

smooth (pseudo-)manifolds

(pseudo-)Riemannian manifolds

Lorentzian manifolds

Lorentzian 4-manifolds



GR: smooth manifold and $\tilde{\nabla}g$



topological manifolds

differentiable (pseudo-)manifolds

smooth (pseudo-)manifolds

(pseudo-)Riemannian manifolds

Lorentzian manifolds

Lorentzian 4-manifolds

GR: assume that spacetime is a Lorentzian 4-manifold



GR: smooth manifold and $\tilde{\nabla}g$

from above:

$$\nabla_{\lambda} g_{\mu\nu} = \partial_{\lambda} g_{\mu\nu} - \Gamma_{\mu\lambda}^{\kappa} g_{\kappa\nu} - \Gamma_{\nu\lambda}^{\kappa} g_{\mu\kappa}$$

GR: smooth manifold and $\tilde{\nabla}g$

from above:

$$\nabla_{\lambda} g_{\mu\nu} = \partial_{\lambda} g_{\mu\nu} - \Gamma_{\mu\lambda}^{\kappa} g_{\kappa\nu} - \Gamma_{\nu\lambda}^{\kappa} g_{\mu\kappa}$$

in the tangent space at x , $\exists$ coordinate basis $\vec{e}^{\bar{\mu}}$ with

$$g_{\bar{\mu}\bar{\nu}} = \eta_{\bar{\mu}\bar{\nu}} = \text{diag}(-1, 1, 1, 1) = g^{\bar{\mu}\bar{\nu}}$$

$$\Rightarrow \partial_{\bar{\lambda}} g_{\bar{\mu}\bar{\nu}} = \partial_{\bar{\lambda}} \eta_{\bar{\mu}\bar{\nu}}$$

GR: smooth manifold and $\tilde{\nabla}g$

from above:

$$\nabla_{\lambda} g_{\mu\nu} = \partial_{\lambda} g_{\mu\nu} - \Gamma^{\kappa}_{\mu\lambda} g_{\kappa\nu} - \Gamma^{\kappa}_{\nu\lambda} g_{\mu\kappa}$$

in the tangent space at x , $\exists$ coordinate basis $\vec{e}^{\bar{\mu}}$ with

$$g_{\bar{\mu}\bar{\nu}} = \eta_{\bar{\mu}\bar{\nu}} = \text{diag}(-1, 1, 1, 1) = g^{\bar{\mu}\bar{\nu}}$$

$$\Rightarrow \partial_{\bar{\lambda}} g_{\bar{\mu}\bar{\nu}} = \partial_{\bar{\lambda}} \eta_{\bar{\mu}\bar{\nu}} = 0$$

GR: smooth manifold and $\tilde{\nabla}g$

from above:

$$\nabla_{\lambda} g_{\mu\nu} = \partial_{\lambda} g_{\mu\nu} - \Gamma_{\mu\lambda}^{\kappa} g_{\kappa\nu} - \Gamma_{\nu\lambda}^{\kappa} g_{\mu\kappa}$$

in the tangent space at x , $\exists$ coordinate basis $\vec{e}^{\bar{\mu}}$ with

$$g_{\bar{\mu}\bar{\nu}} = \eta_{\bar{\mu}\bar{\nu}} = \text{diag}(-1, 1, 1, 1) = g^{\bar{\mu}\bar{\nu}}$$

$$\Rightarrow \partial_{\bar{\lambda}} g_{\bar{\mu}\bar{\nu}} = \partial_{\bar{\lambda}} \eta_{\bar{\mu}\bar{\nu}} = 0$$

$$\text{also, } \Gamma_{\bar{\nu}\bar{\mu}}^{\bar{\lambda}} \vec{e}_{\bar{\lambda}} := \vec{e}_{\bar{\nu},\bar{\mu}} = \partial_{\bar{\mu}} \vec{e}_{\bar{\nu}}$$

GR: smooth manifold and $\tilde{\nabla}g$

from above:

$$\nabla_{\lambda} g_{\mu\nu} = \partial_{\lambda} g_{\mu\nu} - \Gamma^{\kappa}_{\mu\lambda} g_{\kappa\nu} - \Gamma^{\kappa}_{\nu\lambda} g_{\mu\kappa}$$

in the tangent space at x , $\exists$ coordinate basis $\vec{e}^{\bar{\mu}}$ with

$$g_{\bar{\mu}\bar{\nu}} = \eta_{\bar{\mu}\bar{\nu}} = \text{diag}(-1, 1, 1, 1) = g^{\bar{\mu}\bar{\nu}}$$

$$\Rightarrow \partial_{\bar{\lambda}} g_{\bar{\mu}\bar{\nu}} = \partial_{\bar{\lambda}} \eta_{\bar{\mu}\bar{\nu}} = 0$$

$$\text{also, } \Gamma^{\bar{\lambda}}_{\bar{\nu}\bar{\mu}} \vec{e}_{\bar{\lambda}} := \vec{e}_{\bar{\nu},\bar{\mu}} = \partial_{\bar{\mu}} \vec{e}_{\bar{\nu}}$$

but in a Cartesian or Minkowski (vector) space, the basis vectors always point in the same direction and their lengths are fixed

GR: smooth manifold and $\tilde{\nabla}g$

from above:

$$\nabla_{\lambda} g_{\mu\nu} = \partial_{\lambda} g_{\mu\nu} - \Gamma_{\mu\lambda}^{\kappa} g_{\kappa\nu} - \Gamma_{\nu\lambda}^{\kappa} g_{\mu\kappa}$$

in the tangent space at x , $\exists$ coordinate basis $\vec{e}^{\bar{\mu}}$ with

$$g_{\bar{\mu}\bar{\nu}} = \eta_{\bar{\mu}\bar{\nu}} = \text{diag}(-1, 1, 1, 1) = g^{\bar{\mu}\bar{\nu}}$$

$$\Rightarrow \partial_{\bar{\lambda}} g_{\bar{\mu}\bar{\nu}} = \partial_{\bar{\lambda}} \eta_{\bar{\mu}\bar{\nu}} = 0$$

$$\text{also, } \Gamma_{\bar{\nu}\bar{\mu}}^{\bar{\lambda}} \vec{e}_{\bar{\lambda}} := \vec{e}_{\bar{\nu},\bar{\mu}} = \partial_{\bar{\mu}} \vec{e}_{\bar{\nu}}$$

$$M^4 \Rightarrow \Gamma_{\bar{\nu}\bar{\mu}}^{\bar{\lambda}} \vec{e}_{\bar{\lambda}} = 0$$

GR: smooth manifold and $\tilde{\nabla}g$

from above:

$$\nabla_{\lambda} g_{\mu\nu} = \partial_{\lambda} g_{\mu\nu} - \Gamma_{\mu\lambda}^{\kappa} g_{\kappa\nu} - \Gamma_{\nu\lambda}^{\kappa} g_{\mu\kappa}$$

in the tangent space at x , $\exists$ coordinate basis $\vec{e}^{\bar{\mu}}$ with

$$g_{\bar{\mu}\bar{\nu}} = \eta_{\bar{\mu}\bar{\nu}} = \text{diag}(-1, 1, 1, 1) = g^{\bar{\mu}\bar{\nu}}$$

$$\Rightarrow \partial_{\bar{\lambda}} g_{\bar{\mu}\bar{\nu}} = \partial_{\bar{\lambda}} \eta_{\bar{\mu}\bar{\nu}} = 0$$

$$\text{also, } \Gamma_{\bar{\nu}\bar{\mu}}^{\bar{\lambda}} \vec{e}_{\bar{\lambda}} := \vec{e}_{\bar{\nu},\bar{\mu}} = \partial_{\bar{\mu}} \vec{e}_{\bar{\nu}}$$

$$\text{so } \nabla_{\bar{\lambda}} g_{\bar{\mu}\bar{\nu}} = 0$$

GR: smooth manifold and $\tilde{\nabla}g$

from above:

$$\nabla_{\lambda} g_{\mu\nu} = \partial_{\lambda} g_{\mu\nu} - \Gamma_{\mu\lambda}^{\kappa} g_{\kappa\nu} - \Gamma_{\nu\lambda}^{\kappa} g_{\mu\kappa}$$

in the tangent space at x , $\exists$ coordinate basis $\vec{e}^{\bar{\mu}}$ with

$$g_{\bar{\mu}\bar{\nu}} = \eta_{\bar{\mu}\bar{\nu}} = \text{diag}(-1, 1, 1, 1) = g^{\bar{\mu}\bar{\nu}}$$

$$\Rightarrow \partial_{\bar{\lambda}} g_{\bar{\mu}\bar{\nu}} = \partial_{\bar{\lambda}} \eta_{\bar{\mu}\bar{\nu}} = 0$$

$$\text{also, } \Gamma_{\bar{\nu}\bar{\mu}}^{\bar{\lambda}} \vec{e}_{\bar{\lambda}} := \vec{e}_{\bar{\nu},\bar{\mu}} = \partial_{\bar{\mu}} \vec{e}_{\bar{\nu}}$$

$$\text{so } \nabla_{\bar{\lambda}} g_{\bar{\mu}\bar{\nu}} = 0$$

so $\tilde{\nabla}g = 0$ (also $\tilde{\nabla}g^{-1} = 0$) on the tangent space, since if true in one coord system, also true in others

GR: smooth manifold and $\tilde{\nabla}g$

from above:

$$\nabla_{\lambda} g_{\mu\nu} = \partial_{\lambda} g_{\mu\nu} - \Gamma^{\kappa}_{\mu\lambda} g_{\kappa\nu} - \Gamma^{\kappa}_{\nu\lambda} g_{\mu\kappa}$$

in the tangent space at x , $\exists$ coordinate basis $\vec{e}^{\bar{\mu}}$ with

$$g_{\bar{\mu}\bar{\nu}} = \eta_{\bar{\mu}\bar{\nu}} = \text{diag}(-1, 1, 1, 1) = g^{\bar{\mu}\bar{\nu}}$$

$$\Rightarrow \partial_{\bar{\lambda}} g_{\bar{\mu}\bar{\nu}} = \partial_{\bar{\lambda}} \eta_{\bar{\mu}\bar{\nu}} = 0$$

$$\text{also, } \Gamma^{\bar{\lambda}}_{\bar{\nu}\bar{\mu}} \vec{e}_{\bar{\lambda}} := \vec{e}_{\bar{\nu},\bar{\mu}} = \partial_{\bar{\mu}} \vec{e}_{\bar{\nu}}$$

$$\text{so } \nabla_{\bar{\lambda}} g_{\bar{\mu}\bar{\nu}} = 0$$

$$\text{so } \tilde{\nabla}g = 0 = \tilde{\nabla}g^{-1} \text{ on tangent space}$$

$$\dots \tilde{\nabla}g = 0 = \tilde{\nabla}g^{-1} \text{ on } M$$

GR: smooth manifold and $\tilde{\nabla}g$

from above:

$$\nabla_{\lambda} g_{\mu\nu} = \partial_{\lambda} g_{\mu\nu} - \Gamma_{\mu\lambda}^{\kappa} g_{\kappa\nu} - \Gamma_{\nu\lambda}^{\kappa} g_{\mu\kappa}$$

in the tangent space at x , $\exists$ coordinate basis $\vec{e}^{\bar{\mu}}$ with

$$g_{\bar{\mu}\bar{\nu}} = \eta_{\bar{\mu}\bar{\nu}} = \text{diag}(-1, 1, 1, 1) = g^{\bar{\mu}\bar{\nu}}$$

$$\Rightarrow \partial_{\bar{\lambda}} g_{\bar{\mu}\bar{\nu}} = \partial_{\bar{\lambda}} \eta_{\bar{\mu}\bar{\nu}} = 0$$

$$\text{also, } \Gamma_{\bar{\nu}\bar{\mu}}^{\bar{\lambda}} \vec{e}_{\bar{\lambda}} := \vec{e}_{\bar{\nu},\bar{\mu}} = \partial_{\bar{\mu}} \vec{e}_{\bar{\nu}}$$

$$\text{so } \nabla_{\bar{\lambda}} g_{\bar{\mu}\bar{\nu}} = 0$$

$$\text{so } \tilde{\nabla}g = 0 = \tilde{\nabla}g^{-1} \text{ on tangent space}$$

$$\dots \tilde{\nabla}g = 0 = \tilde{\nabla}g^{-1} \text{ on } M$$

$$\dots \Gamma_{\mu\nu}^{\lambda} = \Gamma_{\nu\mu}^{\lambda} \text{ in any coord. basis (symmetric defn)}$$

GR: smooth manifold and $\tilde{\nabla}g$

from above:

$$\nabla_{\lambda} g_{\mu\nu} = \partial_{\lambda} g_{\mu\nu} - \Gamma^{\kappa}_{\mu\lambda} g_{\kappa\nu} - \Gamma^{\kappa}_{\nu\lambda} g_{\mu\kappa}$$

in the tangent space at x , $\exists$ coordinate basis $\vec{e}^{\bar{\mu}}$ with

$$g_{\bar{\mu}\bar{\nu}} = \eta_{\bar{\mu}\bar{\nu}} = \text{diag}(-1, 1, 1, 1) = g^{\bar{\mu}\bar{\nu}}$$

$$\Rightarrow \partial_{\bar{\lambda}} g_{\bar{\mu}\bar{\nu}} = \partial_{\bar{\lambda}} \eta_{\bar{\mu}\bar{\nu}} = 0$$

$$\text{also, } \Gamma^{\bar{\lambda}}_{\bar{\nu}\bar{\mu}} \vec{e}_{\bar{\lambda}} := \vec{e}_{\bar{\nu},\bar{\mu}} = \partial_{\bar{\mu}} \vec{e}_{\bar{\nu}}$$

$$\text{so } \nabla_{\bar{\lambda}} g_{\bar{\mu}\bar{\nu}} = 0$$

$$\text{so } \tilde{\nabla}g = 0 = \tilde{\nabla}g^{-1} \text{ on tangent space}$$

$$\dots \tilde{\nabla}g = 0 = \tilde{\nabla}g^{-1} \text{ on } M$$

$$\dots \Gamma^{\lambda}_{\mu\nu} = \Gamma^{\lambda}_{\nu\mu} \text{ in any coord. basis (symmetric defn)}$$

...

$$\Gamma^{\lambda}_{\mu\nu} = \frac{1}{2} g^{\lambda\kappa} (\partial_{\mu} g_{\nu\kappa} + \partial_{\nu} g_{\mu\kappa} - \partial_{\kappa} g_{\mu\nu}) \text{ in a coordinate basis}$$

GR:

w:Einstein field equations

GR:

w:Einstein field equations

GR:

w:Einstein field equations

GR:

w:Einstein field equations

GR:

w:Einstein field equations

GR:

w:Einstein field equations

GR:

w:Einstein field equations

GR:

w:Einstein field equations

GR:

w:Equivalence principle

can be thought of as a *consequence* of the model

GR:

w:Equivalence principle

can be thought of as a *consequence* of the model

GR:

w:Equivalence principle

can be thought of as a *consequence* of the model

GR:

w:Equivalence principle

can be thought of as a *consequence* of the model

GR:

w:Equivalence principle

can be thought of as a *consequence* of the model

GR:

w:Equivalence principle

can be thought of as a *consequence* of the model

GR:



w:Equivalence principle

can be thought of as a *consequence* of the model



GR:



w:Equivalence principle

can be thought of as a *consequence* of the model



GR:

w:Schwarzschild metric

GR:

w:Schwarzschild metric

GR:

w:Schwarzschild metric

GR:

w:Schwarzschild metric

GR:

w:Schwarzschild metric

GR:

w:Schwarzschild metric

GR:

w:Schwarzschild metric

GR:

w:Schwarzschild metric

GR:



w:Friedmann-Lemaître-Robertson-Walker metric



GR:



w:Friedmann-Lemaître-Robertson-Walker metric



GR:



w:Friedmann-Lemaître-Robertson-Walker metric



GR:



w:Friedmann-Lemaître-Robertson-Walker metric



GR:



w:Friedmann-Lemaître-Robertson-Walker metric



GR:



w:Friedmann-Lemaître-Robertson-Walker metric



GR:



w:Friedmann-Lemaître-Robertson-Walker metric



GR:



w:Friedmann-Lemaître-Robertson-Walker metric



GR: maxima

maxima - component tensor packet ctensor; itensor

GR: maxima

maxima - component tensor packet ctensor; itensor

GR: maxima

maxima - component tensor packet ctensor; itensor

GR: maxima

maxima - component tensor packet ctensor; itensor

GR: maxima

maxima - component tensor packet ctensor; itensor

GR: maxima

maxima - component tensor packet ctensor; itensor

GR: maxima

maxima - component tensor packet ctensor; itensor

GR: maxima

maxima - component tensor packet ctensor; itensor

GR: an approximation method: ADM

+ w:ADM formalism



GR: an approximation method: ADM

+ w:ADM formalism



GR: an approximation method: ADM

+ w:ADM formalism



GR: an approximation method: ADM

+ w:ADM formalism



GR: an approximation method: ADM

+ w:ADM formalism



GR: an approximation method: ADM

+ w:ADM formalism



GR: an approximation method: ADM

+ w:ADM formalism



GR: an approximation method: ADM

+ w:ADM formalism



GR: a numerical method: cactus



+ Cactus - <http://cactuscode.org>



GR: a numerical method: cactus



+ Cactus - <http://cactuscode.org>



GR: a numerical method: cactus



+ Cactus - <http://cactuscode.org>



GR: a numerical method: cactus



+ Cactus - <http://cactuscode.org>



GR: a numerical method: cactus



+ Cactus - <http://cactuscode.org>



GR: a numerical method: cactus



+ Cactus - <http://cactuscode.org>



GR: a numerical method: cactus



+ Cactus - <http://cactuscode.org>



GR: a numerical method: cactus

+ Cactus - <http://cactuscode.org>